

# Section 5. Risk Assessment

## 5.6 Flood

For the 2014 Plan update, the hazard profile and vulnerability assessment were significantly enhanced to reflect updated, best available data. A recap of each Federal Emergency Management Agency (FEMA) major disaster or emergency declaration event has been provided, along with events that did not result in a declaration, when available. To assess vulnerability, the latest FEMA mapping was used, including preliminary work maps released in 2013 for coastal areas. The vulnerability assessment also includes a comparison between historic and current National Flood Insurance Program (NFIP) data to demonstrate the changes that have occurred within the State over time. This information can be used by both the state agencies in developing mitigation strategies, as well as the local jurisdictions as they develop their mitigation plans.

## 5.6.1 Profile

This section provides general information on the flood hazard which includes riverine (inland) flooding, coastal flooding, ice jams, stormwater flooding, and tsunamis. Flooding caused by dam and levee failure is discussed in Section 5.3 (Dam and Levee Failure), and storm surge is discussed in Section 5.8 (Hurricane and Tropical Storms).

## **Hazard Description**

Floods are one of the most common natural hazards in the United States They can develop slowly over a period of days or develop quickly, with disastrous effects that can be local (impacting a neighborhood or community) or regional (affecting entire river basins, coastlines and multiple counties or states) (FEMA 2008). Most communities in the United States have experienced some kind of flooding after spring rains, heavy thunderstorms, coastal storms, or winter snow thaws (George Washington University 2001). Floods are frequent and costly natural hazards in New Jersey in terms of human hardship and economic loss, particularly to communities that lie within flood-prone areas or floodplains of a major water source.

The flood-related hazards most likely to affect New Jersey are riverine (inland) flooding and coastal flooding. Other flood-related hazards that have historically occurred and will continue to affect the State include: flooding associated with ice jams, flooding associated with tsunamis, stormwater flooding due to local drainage and high groundwater levels, and storm surge/coastal flooding. Each is described below, along with the sub-categories associated with each hazard type. Storm surge and coastal flooding are discussed further in Section 5.8 (Hurricanes and Tropical Storms).

## Riverine (Inland) Flooding

Riverine floods are the most common flood type. They occur along a channel and include overbank and flash flooding. Channels are defined, ground features that carry water through and out of a watershed. They may be called rivers, creeks, streams, or ditches. When a channel receives too much water, the excess water flows over its banks and inundates low-lying areas (FEMA 2008; The Illinois Association for Floodplain and Stormwater Management 2006).


Flash floods are "a rapid and extreme flow of high water into a normally dry area, or a rapid water level rise in a stream or creek above a predetermined flood level, beginning within six hours of the causative event (e.g., intense rainfall, dam failure, ice jam). However, the actual time threshold may vary in different parts of the



country. Ongoing flooding can intensify to flash flooding in cases where intense rainfall results in a rapid surge of rising flood waters" (National Weather Service [NWS] 2009).

A floodplain is defined as the land adjoining the channel of a river, stream, ocean, lake, or other watercourse or water body that becomes inundated with water during a flood. Most often floodplains are referred to as 100-year floodplains. A 100-year floodplain is not a flood that will occur once every 100 years, rather it is a flood that has a one-percent chance of being equaled or exceeded each year. Thus, the 100-year flood could occur more than once in a relatively short period of time. Due to this misleading term, FEMA has properly defined it as the one-percent annual chance flood. This one-percent annual chance flood is now the standard used by most federal and state agencies and by the NFIP (FEMA 2002).

Figure 5.6-1 depicts the flood hazard area, the flood fringe, and the floodway areas of a floodplain.



#### Figure 5.6-1. Floodplain

Source: New Jersey Department of Environmental Protection (NJDEP) 2009

## **Coastal Flooding**

Coastal flooding generally occurs along the coasts of oceans, bays, estuaries, coastal rivers, and large lakes. Coastal floods are the submersion of land areas along the ocean coast and other inland waters caused by seawater over and above normal tide action. Coastal flooding is a result of the storm surge where local sea levels rise often resulting in weakened or destroyed coastal structures. Hurricanes and tropical storms, severe storms, and Nor'Easters cause most of the coastal flooding in New Jersey. Coastal flooding has many of the same problems identified for riverine flooding but also has additional problems such as beach erosion; loss or submergence of wetlands and other coastal ecosystems; saltwater intrusion; high water tables; loss of coastal recreation areas, beaches, protective sand dunes, parks, and open space; and loss of coastal structures. Coastal structures can include sea walls, piers, bulkheads, bridges, or buildings (FEMA 2011).

There are several forces that occur with coastal flooding:

- *Hydrostatic forces* against a structure are created by standing or slowly moving water. Flooding can cause vertical hydrostatic forces, or flotation. These types of forces are one of the main causes of flood damage.
- *Hydrodynamic forces* on buildings are created when coastal floodwaters move at high velocities. These high-velocity flows are capable of destroying solid walls and dislodging buildings with inadequate foundations. High-velocity flows can also move large quantities of sediment and debris



that can cause additional damage. In coastal areas, high-velocity flows are typically associated with one or more of the following:


- Storm surge and wave run-up flowing landward through breaks in sand dunes or across low-lying areas
- o Tsunamis
- $\circ$   $\;$  Outflow of floodwaters driven into bay or upland areas
- Strong currents parallel to the shoreline, driven by waves produced from a storm
- High-velocity flows

High-velocity flows can be created or exacerbated by the presence of manmade or natural obstructions along the shoreline and by weak points formed by roads and access paths that cross dunes, bridges or canals, channels, or drainage features.

- *Waves* can affect coastal buildings from breaking waves, wave run-up, wave reflection and deflection, and wave uplift. The most severe damage is caused by breaking waves. The force created by these types of waves breaking against a vertical surface is often at least 10 times higher than the force created by high winds during a coastal storm.
- *Flood-borne debris* produced by coastal flooding events and storms typically includes decks, steps, ramps, breakaway wall panels, portions of or entire houses, heating oil and propane tanks, cars, boats, decks and pilings from piers, fences, erosion control structures, and many other types of smaller objects. Debris from floods are capable of destroying unreinforced masonry walls, light wood-frame construction, and small-diameter posts and piles (FEMA 2011).

According to the 2011 Coastal Construction Manual, FEMA P-55, Zone V (including Zones VE, V1-30, and V) identifies the Coastal High Hazard Area. This is the portion of the special flood hazard area (SFHA) that extends from offshore to the inland limit of a primary frontal dune along an open coast and any other portion of the SFHA that is subject to high-velocity wave action from storms or seismic sources. The boundary of Zone V is generally based on wave heights (3 feet or greater) or wave run-up depths (3 feet or greater). Zone V can also be mapped based on the wave overtopping rate (when waves run up and over a dune or barrier). Zone A or AE, identify portions of the SFHA that are not within the Coastal High Hazard Area. These zones are used to designate both coastal and non-coastal SFHAs. Regulatory requirements of the NFIP for buildings located in Zone A are the same for both coastal and riverine flooding hazards. Zone AE in coastal areas is divided by the limit of moderate wave action (LiMWA). The LiMWA represents the landward limit of the 1.5-foot wave (FEMA 2011).

The area between the LiMWA and the Zone V limit is known as the Coastal A-zone (for building codes and standard purposes) and as the Moderate Wave Action area (by FEMA flood mappers). This area is subject to wave heights between 1.5 and 3 feet during the base flood. The area between the LiMWA and the landward limit of Zone A is known as the Minimal Wave Action area, and is subject to wave heights less than 1.5 feet during the base flood (FEMA P-55 2011). Figure 5.6-2 shows a typical transect illustrating Zone V, the Coastal A-zone and Zone A, and the effects of energy dissipation and regeneration of a wave as it moves inland. Wave elevations are decreased by obstructions such as vegetation and rising ground elevation (FEMA 2011).



#### Figure 5.6-2. Transect Schematic of Zone V, Coastal A-zone, and Zone A

#### Ice Jam

As per the Northeast States Emergency Consortium and FEMA, an ice jam is an accumulation of ice that acts as a natural dam and restricts flow of a body of water. Ice jams occur when warm temperatures and heavy rains cause rapid snowmelt. The melting snow, combined with the heavy rain, causes frozen rivers to swell. The rising water breaks the ice layers into large chunks, which float downstream and often pile up near narrow passages and obstructions (bridges and dams). Ice jams may build up to a thickness great enough to raise the water level and cause flooding (FEMA 2008). Ice jams may also be caused by frazil ice, which forms when mist freezes then floats down a river, stream, or creek.

There are two different types of ice jams: freeze-up and breakup. Freeze-up jams occur in the early to midwinter when floating ice may slow or stop due to a change in water slope as it reaches an obstruction to movement. Breakup jams occur during periods of thaw, generally in late winter and early spring. The ice cover breakup is usually associated with a rapid increase in runoff and corresponding river discharge due to a heavy rainfall, snowmelt, or warmer temperatures (United States Army Corps of Engineers [USACE] 2002).

## Stormwater Flooding

Stormwater flooding described below is due to local drainage issues and high groundwater levels. Locally, heavy precipitation may produce flooding in areas other than delineated floodplains or along recognizable channels. If local conditions cannot accommodate intense precipitation through a combination of infiltration and surface runoff, water may accumulate and cause flooding problems. During winter and spring, frozen ground and snow accumulations may contribute to inadequate drainage and localized ponding. Flooding issues of this nature generally occur in areas with flat gradients and generally increase with urbanization which speeds the accumulation of floodwaters because of impervious areas. Shallow street flooding can occur unless channels have been improved to account for increased flows (FEMA 1997).



High groundwater levels can be a concern and cause problems even where there is no surface flooding. Basements are susceptible to high groundwater levels. Seasonally high groundwater is common in many areas, while elsewhere high groundwater occurs only after a long periods of above-average precipitation (FEMA 1997).

Urban drainage flooding is caused by increased water runoff due to urban development and drainage systems. Drainage systems are designed to remove surface water from developed areas as quickly as possible to prevent localized flooding on streets and other urban areas. They make use of a closed conveyance system that channels water away from an urban area to surrounding streams. This bypasses the natural processes of water filtration through the ground, containment, and evaporation of excess water. Since drainage systems reduce the amount of time the surface water takes to reach surrounding streams, flooding in those streams can occur more quickly and reach greater depths than prior to development in that area (FEMA 2008).

#### Tsunami

FEMA and NOAA state that tsunamis are a series of traveling ocean waves created by sudden displacements of the ocean floor (earthquakes), landslides, or volcanic activity. A tsunami can move hundreds of miles per hour in the open ocean and crash into land with waves exceeding 100 feet in height (FEMA 2009).

A tsunami consists of a series of high-energy waves that travel outward, like pond ripples, from the area where the tsunami originated. The sequence of tsunami waves arrives at the shoreline over an extended period of time and build height as it get closer (FEMA 2007; Humboldt County Hazard Mitigation Plan 2008). A tsunami approaching the shoreline may take three forms:

- Non-breaking waves that act as a rapidly rising tide
- A large, turbulent wall-like wave (bore)
- A series of partially developed waves

#### Source: Humboldt County Hazard Mitigation Plan 2008

There are two types of tsunamis: local and distant. A locally generated tsunami is caused by an undersea disturbance near the coast. They have minimal warning times and may be accompanied by earthquake damage due to ground shaking, surface faulting, liquefaction, or landslides. A local tsunami, due to its close proximity to the coast, leaves few options for escaping, except to run to high ground. Distant tsunamis may travel for hours before striking a coastline, leaving enough time for warning (Humboldt County Hazard Mitigation Plan 2008; Grays Harbor County Hazard Mitigation Plan 2005).

## Federal Flood Programs

#### National Flood Insurance Program

The NFIP makes federally backed flood insurance available to homeowners, renters, and business owners in participating communities. For most participating communities, FEMA has prepared a detailed Flood Insurance Study (FIS). The study presents water surface elevations for floods of various magnitudes, including the 1% annual chance flood and the 0.2% annual chance flood (the 500-year flood). Base flood elevations and the boundaries of the 100- floodplains are shown on Flood Insurance Rate Maps (FIRMs), which are the principal tool for identifying the extent and location of the flood hazard.

The FIRMs depict SFHAs - those areas subject to inundation from the 1% annual chance flood (also known as the Base Flood or the 100-Year Flood). Those areas are defined as follows:



- Zones A1-30 and AE: SFHAs that are subject to inundation by the base flood, determined using detailed hydraulic analysis. Base Flood Elevations are shown within these zones.
- Zone A (Also known as Unnumbered A-zones): SFHAs where no Base Flood Elevations or depths are shown because detailed hydraulic analyses have not been performed,.
- Zone AO: SFHAs subject to inundation by types of shallow flooding where average depths are between one and three feet. These are normally areas prone to shallow sheet flow flooding on sloping terrain.
- Zone VE, V1-30: SFHAs along coasts that are subject to inundation by the base flood with additional hazards due to waves with heights of three feet or greater. Base Flood Elevations derived from detailed hydraulic analysis are shown within these zones.
- Zone B and X (shaded): Zones where the land elevation as been determined to be above the Base Flood Elevation, but below the 500-year flood elevation. These zones are not SFHAs.
- Zones C and X (unshaded): Zones where the land elevation has been determined to be above both the Base Flood Elevation and the 500-year flood elevation. These zones are not SFHAs.

There are approximately 245,806 NFIP policies in New Jersey. Of those policies, 16,017 are considered repetitive loss (RL) and 2,097 are considered severe repetitive loss (SRL). To be eligible for the NFIP, certain criteria must be met and claim payments must have occurred within 10 years of each other. If there are multiple losses at the same location within 10 days of each other, these claims are counted within one loss.

## Flood Insurance Studies (FIS)

In addition to FIRM and Digital Flood Insurance Rate Maps (DFIRM), FEMA also provides FISs for entire counties and individual jurisdictions. These studies aid in the administration of the National Flood Insurance Act of 1968 and the Flood Disaster Protection Act of 1973. They are narrative reports of countywide flood hazards, including descriptions of the flood areas studied, the engineered methods used, principal flood problems, flood protection measures, and graphic profiles of the flood sources.

#### Risk Mapping, Assessment, and Planning (Risk MAP)

Risk MAP is a FEMA program that provides communities with flood information and tools to enhance their mitigation plans and take action to protect their citizens. It builds on flood hazard data and maps produced during the Flood Map Modernization (Map Mod) program. Through more precise flood mapping products, risk assessment tools, and planning and outreach support, Risk MAP strengthens local ability to make informed decisions about reducing risk. It combines quality engineering with state-of-the-art flood hazard data to assist communities in planning and preventing risk using the most current information.

Risk MAP collaborates with state, tribal, and local governments and delivers quality data that increases public awareness and leads to action that reduces risk to property and life. Risk MAP focuses on products and services beyond the traditional FIRMs and works with officials to help put flood risk data and assessment tools to use. Risk MAP also helps effectively communicate risk to citizens and enable communities to enhance their mitigation plans and actions (FEMA 2012).

The goals of Risk MAP are as follows:

- Flood Hazard Data addresses gaps in flood hazard data to form a solid foundation for risk assessment, floodplain management, and actuarial soundness of the NFIP.
- Public Awareness/Outreach ensures that a measurable increase of the public's awareness and understanding of risk results in a measurable reduction of current and future vulnerability.



- Hazard Mitigation Planning leads and supports states, local, and tribal communities to effectively engage in risk-based mitigation planning resulting in sustainable actions that reduce or eliminate risks to life and property from natural hazards.
- Enhanced Digital Platform provides an enhanced digital platform that improves management of Risk MAP, conserves information produced by Risk MAP, and improves communication and sharing of risk data and related products to all levels of government and the public.
- Alignment and Synergies aligns risk analysis programs and develops synergies to enhance decisionmaking capabilities through effective risk communication and management.

FEMA headquarters and regional offices lead a team of contractors and stakeholders to deliver its Risk MAP program. The team is made up of the following:

- FEMA Headquarters responsible for overall program implementation
- FEMA Regions manage regional flood map production and help implement the Risk MAP outreach strategy
- State, Local, and Tribal entities help ensure that updated mapping information is used to make informed decisions regarding risk
- Program Management Contractor provide general oversight for Risk MAP including integration of activities, development and implementation of a national outreach strategy, and stakeholder relations
- Production and Technical Services Contractors update flood hazard data and maps
- Customer and Data Services Contractor provide the digital platform for sharing flood mapping products and information

The NJDEP executed a Cooperating Technical Partners (CTP) partnership agreement with FEMA on May 16, 2006. Since that time, NJDEP has become a full CTP partner with FEMA. Under the CTP agreement, the NJDEP works as a contractor to FEMA Region II on the production of regulatory and non-regulatory Flood Risk MAP products for the State of New Jersey.

#### Biggert-Water Flood Insurance Reform Act of 2012

In July 2012, the United States Congress passed the Biggert-Water Flood Insurance Reform Act of 2012 (BW-12) which calls on FEMA and other agencies to make a number of changes to the way the NFIP is run. Key provisions of the legislation will require the NFIP to raise rates to reflect true flood risk, make the program more financially stable, and change how FIRM updates impact policyholders. BW-12 also eliminated the Repetitive Flood Claims and Severe Repetitive Loss programs and made significant changes to the Flood Mitigation Assistance (FMA) program. These changes include:

- The definitions of repetitive loss and severe repetitive loss properties were modified and are as follows:
  - A severe repetitive loss property is a structure that is covered under a contract for flood insurance made available under the NFIP. These properties have incurred flood-related damage for which four or more separate claims payments have been made under flood insurance coverage with the amount of each such claim exceeding \$5,000 and with the cumulative amount of such claims payments exceeding \$20,000. Or for which at least two separate claims payments have been made under such coverage, with the cumulative amount of such claims exceeding that market value of the insured structure.
  - A repetitive loss property is a structure covered by a contract for flood insurance made available under the NFIP that has incurred flood-related damage on two occasions, in which the cost of the repair, on average, equaled or exceeded 25% of market value of the structure at



the time of each such flood event. Also, at the time of the second incidence of flood-related damage, the contract for flood insurance contains increased cost of compliance coverage.


- There is no longer a state cap of \$10 million or a community cap of \$3.3 million for any five-year period
- There is no longer a limit on in-kind contributions for the non-federal cost share
- Mitigation reconstruction is an eligible activity
- Cost-share requirements have changed to allow more federal funds for properties with repetitive flood claims and severe repetitive loss properties
- The development or update of mitigation plans shall not exceed \$50,000 federal share to any applicant or \$25,000 federal share to any subapplicant
- There is no longer a restriction that a planning grant can be awarded not more than once every five years to a state or community (FEMA 2013a)

Homeowners of certain older properties in high-risk areas had been charged premiums that do not reflect the full flood risk. Only properties known as "pre-FIRM" were eligible for these subsidies. Although only approximately 20% of NFIP policies nationwide are subsidized, 37.1% of New Jersey policies are considered to be "pre-FIRM". BW-12 requires FEMA to phase out these subsidies for certain properties and prohibits FEMA from offering subsidies for other pre-FIRM properties. Not all subsidies will be removed the same way at the same time (FEMA 2013a). Increases to pre-FIRM subsided rates include the following:

- Owners of non-primary residences with pre-FIRM subsidized rates are scheduled to see a 25% annual increase until full-risk rates are reached, unless superseded by pending Congressional legislation.
- By October 1, 2013:
  - Owners of businesses with pre-FIRM subsidized rates will see a 25% annual increase until full-risk rates are reached
  - Owners of properties of one to four residences with a pre-FIRM subsidized rate that have experienced severe or repetitive flooding will see a 25% annual increase until full-risk rates are reached
  - Pre-FIRM subsidized policies first in effect on or after July 6, 2012 will move directly to fullrisk rates
  - Pre-FIRM subsidized policies on homes purchased on or after July 6, 2012 will move directly to full-risk rates
  - Lapsed pre-FIRM subsidized policies reinstated on or after October 4, 2012 will move directly to full-risk rates (FEMA 2013a)

Upon a revised or updated flood map, BW-12 requires adjustment and phase-in rates over five years to accurately reflect the current risk of flood to properties (FEMA 2013a).

In New Jersey there are a total of 2,097 SRL properties. The county with the highest number of SRL properties is Passaic County (610 SRL properties). Figure 5.6-3 shows the number of severe repetitive loss properties in New Jersey.



#### Figure 5.6-3. Number of Severe Repetitive Loss Properties per County in New Jersey

Source: NJDEP 2013 Note: Data current as of June 2013.



## Community Rating System (CRS) Program

The CRS is a voluntary program within the NFIP encouraging floodplain management activities that exceed the minimum NFIP requirements. Flood insurance premiums are discounted to reflect the reduced flood risk to meet the CRS goals of reducing flood losses, facilitating accurate insurance rating, and promoting awareness of flood insurance in the community.

For participating communities, flood insurance premium rates are discounted in increments of 5%. For example, a Class 1 community receives a 45% premium discount, and a Class 9 community receives a 5% discount. Class 10 communities do not participate in the CRS and therefore do not receive a discount. The CRS classes for local communities are based on 18 creditable activities in the following categories:

- Public information
- Mapping and regulations
- Flood damage reduction
- Flood preparedness

CRS activities (discussed below) can help save lives and reduce property damage. Communities participating in the CRS represent a significant portion of the nation's flood risk; over 66% of the NFIP's policy base is located in these communities. Small and large communities participate in and receive premium discounts through the CRS. These communities represent a mixture of flood risks, including both coastal and riverine flood risks. The Insurance Services Office (ISO) administers the CRS program under contract to FEMA.

As of October 2013, there were 82 communities within the State of New Jersey participating in the CRS program. The participating communities are shown in Table 5.6-1. These communities represent 0.03% of the flood insurance policy base within the State. The CRS classifications in New Jersey range from a high of Class 10 (no discount) to a low of Class 5 (25% discount). The New Jersey Dam Safety program, state stormwater management requirements, and the development of all hazard mitigation plans are some of the efforts at the state level that provide CRS credits for all New Jersey municipalities. Communities are encouraged to adopt freeboard elevation requirements, which also provide CRS credits. Many municipalities in New Jersey are small and lack the professional support to fill out a CRS application, or do not have the flood insurance policy base to make it worthwhile. However, Community Assistance Visits (CAV), Community Assistance Contacts (CAC), technical assistance contacts, and workshops help to promote the CRS program in these small towns.

As of October 2013, New Jersey has:

- 20 communities with a Class 10 (0%) premium reduction;
- 10 communities with a Class 9 rating (5% premium reduction);
- 18 communities with a Class 8 rating (10% premium reduction);
- 12 communities with a Class 7 rating (15% premium reduction);
- 13 communities with a Class 6 rating (20% premium reduction); and
- 9 communities with a Class 5 rating (25% premium reduction).

The total annual flood insurance premium CRS discount for the State as of February 25, 2014 was \$25,447,131. This represents 10.6% of the total annual premium (\$240,939,675) for the State.

Of the participating CRS communities, 20 of them had their CRS classifications rescinded due to failure to meet annual participation requirements. These communities are receiving no CRS Discount.



## Table 5.6-1. Participating CRS Communities in New Jersey

Community #	Community	CRS Entry Date	Current Effective Date	Current Class	% Discount for SFHA	% Discount for Non- SFHA	Status
340312	Aberdeen, Township of	5/1/2010	5/1/2010	9	5	5	C
345278	Atlantic City, City of	10/1/1992	10/1/2000	9	5	5	C
345279	Avalon, Borough of	10/1/1996	10/1/2013	5	25	10	C
340396	Barnegat, Township of	10/1/1992	10/1/1997	10	0	0	R
345280	Barnegat Light, Borough of	10/1/1992	10/1/2001	8	10	5	C
345281	Bay Head, Borough of	10/1/1993	10/1/2013	6	20	10	C
345282	Beach Haven, Borough of	10/1/1991	10/1/2013	5	25	10	C
340427	Bedminster, Township of	10/1/1996	5/1/2007	6	20	10	C
340369	Berkeley, Township of	10/1/1992	10/1/2013	6	20	10	C
340459	Berkeley Heights, Township of	10/1/1994	4/1/1999	10	0	0	R
340428	Bernards, Township of	10/1/2010	10/1/2010	8	10	5	C
340178	Bloomfield, Township of	10/1/1992	10/1/1997	10	0	0	R
340289	Bradley Beach, Borough of	10/1/1995	10/1/2000	7	15	5	C
345286	Brigantine, City of	10/1/1992	10/1/2009	6	20	10	C
345287	Burlington, City of	4/1/1998	10/1/2003	8	10	5	C
345288	Cape May City, City of	10/1/1994	10/1/2013	6	20	10	C
345289	Cape May Point, Borough of	10/1/1993	10/1/2013	6	20	10	C
345292	Denville, Township of	10/1/2011	10/1/2011	9	5	5	C
340031	Englewood, City of	10/1/1991	10/1/2001	10	0	0	R
345295	Fairfield, Township of	5/1/2013	5/1/2013	6	20	10	C
340434	Franklin, Township of	5/1/2010	5/1/2010	7	15	5	C
340037	Garfield, City of	5/1/2012	5/1/2012	9	5	5	С
340204	Greenwich, Township of	5/1/2007	5/1/2007	9	5	5	C
340246	Hamilton, Township of	10/1/1992	10/1/2002	8	10	5	С
345296	Harvey Cedars, Borough of	10/1/1991	10/1/1999	8	10	5	С

Page 5.6-11



Community #	Community	CRS Entry Date	Current Effective Date	Current Class	% Discount for SFHA	% Discount for Non- SFHA	Status
340298	Hazlet, Township of	5/1/2011	10/1/2013	6	50	10	С
340376	Lacey, Township of	10/1/1992	10/1/1993	10	0	0	R
340237	Lambertville, City of	5/1/2012	5/1/2012	8	10	5	С
340379	Lavallette, Borough of	5/1/2004	10/1/2013	6	50	10	C
345300	Lincoln Park, Borough of	10/1/1991	10/1/2006	10	0	0	R
340467	Linden, City of	10/1/1991	10/1/2002	8	10	5	C
340401	Little Falls, Township of	5/1/2010	5/1/2010	9	5	5	С
340046	Little Ferry, Borough of	10/1/1993	10/1/1994	10	0	0	R
340047	Lodi, Borough of	10/1/1992	10/1/1993	10	0	0	R
345301	Long Beach, Township of	10/1/1992	10/1/2013	5	25	10	C
345302	Longport, Borough of	10/1/1995	10/1/2013	5	25	10	С
345303	Manasquan, Borough of	10/1/1992	10/1/2009	7	15	5	C
340383	Mantoloking, Borough of	10/1/1992	10/1/2013	5	25	10	С
345304	Margate City, City of	10/1/1992	10/1/2013	5	25	10	C
340313	Middleton Township	5/1/2012	10/1/2013	6	20	10	С
340188	Montclair, Township of	10/1/1994	10/1/1995	10	0	0	R
340517	Mullica, Township of	10/1/1994	5/1/2008	10	0	0	R
340209	National Park, Borough of	10/1/2012	10/1/2012	9	5	5	C
340570	New Jersey Meadowlands Commission	10/1/1992	5/1/2009	7	15	5	С
345307	North Plainfield, Borough of	10/1/1992	10/1/2009	8	10	5	C
345308	North Wildwood, City of	10/1/2000	10/1/2000	7	15	5	C
345309	Oakland, Borough of	10/1/1995	10/1/1996	10	0	0	R
340518	Ocean, Township of	10/1/1995	5/1/2012	10	0	0	R
345310	Ocean City, City of	10/1/1992	10/1/2013	6	20	10	C
340320	Oceanport, Borough of	5/1/2010	5/1/2010	8	10	5	С
340110	Palmyra, Borough of	10/1/2009	10/1/2009	8	10	5	C



Community #	Community	CRS Entry Date	Current Effective Date	Current Class	% Discount for SFHA	% Discount for Non- SFHA	Status
340355	Parsippany-Troy Hills, Township of	10/1/1991	5/1/2009	10	0	0	R
340512	Pennsville, Township of	10/1/1992	10/1/1997	10	0	0	R
345311	Pequannock, Township of	10/1/1991	10/1/2011	7	15	5	С
345312	Plainfield, City of	10/1/1991	10/1/1998	10	0	0	R
345313	Point Pleasant, Borough of	10/1/1993	10/1/2013	8	10	5	C
340388	Point Pleasant Beach, Borough of	10/1/1992	5/1/2009	9	5	5	C
345528	Pompton Lakes, Borough of	10/1/1991	5/1/2013	5	25	10	C
345314	Rahway, City of	10/1/1992	5/1/2013	6	20	10	C
340067	Ridgewood, Village of	10/1/1992	10/1/2002	7	15	5	C
340359	Riverdale, Borough of	10/1/1994	10/1/1994	9	5	5	C
340070	Rochelle Park, Township of	10/1/2006	10/1/2006	8	10	5	C
340472	Roselle, Borough of	10/1/1992	5/1/2013	7	15	5	C
340474	Scotch Plains, Township of	10/1/1994	10/1/1995	10	0	0	R
345317	Sea Bright, Borough of	10/1/1992	10/1/1997	10	0	0	R
345318	Sea Isle City, City of	10/1/1992	10/1/2013	5	25	10	C
345319	Seaside Park, Borough of	10/1/1992	10/1/2006	8	10	5	C
345320	Ship Bottom, Borough of	10/1/1992	5/1/2009	7	15	5	C
340329	Spring Lake, Borough of	10/1/1994	10/1/1999	8	10	5	C
340393	Stafford, Township of	10/1/1991	10/1/2013	5	25	10	C
345323	Stone Harbor, Borough of	10/1/1994	5/1/2009	7	15	5	C
345324	Surf City, Borough of	10/1/1992	10/1/2008	7	15	5	C
345293	Toms River, Township of	10/1/1992	5/1/2013	8	10	5	C
340395	Tuckerton, Borough of	10/1/1993	10/1/1998	10	0	0	R
340331	Union Beach, Borough of	10/1/2003	10/1/2003	8	10	5	C
340159	Upper Township	10/1/2011	10/1/2013	6	20	10	C
345326	Ventnor, City of	10/1/1992	5/1/2010	7	15	5	C



Community #	Community	CRS Entry Date	Current Effective Date	Current Class	% Discount for SFHA	% Discount for Non- SFHA	Status
340446	Warren, Township of	5/1/2010	5/1/2010	9	5	5	С
345327	Wayne, Township of	10/1/1991	10/1/2000	8	10	5	C
345328	West Wildwood, Borough of	10/1/1993	10/1/2005	10	0	0	R
345330	Wildwood Crest, Borough of	10/1/1993	10/1/2003	8	10	5	C
345331	Woodbridge, Township of	10/1/1992	10/1/1997	10	0	0	R

Source: FEMA 2012a

Note: For the purpose of determining CRS discounts, all AR and A99 Zones are treated as non-SFHAs

Zone A99 is the flood insurance rate zone that corresponds to areas of the 1% chance floodplains that will be protected by a Federal flood protection system where construction has reached specified statutory milestones. Mandatory flood insurance purchase requirements apply.

Zone AR is the flood insurance rate zone used to depict areas protected from flood hazards by flood control structures, such as a levee, that are being restored. Mandatory purchase requirements for flood insurance will apply in Zone AR, but the rate will not exceed the rate for unnumbered A zones if the structure is built in compliance with Zone AR floodplain management regulations.

The CRS uses a class rating system that has ratings from 10 to 1. Each CRS Class improvement produces a 5% greater discount on flood insurance premiums for properties in the SFHA, with a Class 1 community receiving the maximum 45% premium reduction.

% percent

C Current

CRS Community rating system

R Rescinded

SFHA Special Flood Hazard Area




#### Location

Flooding in New Jersey is often the direct result of frequent weather events such as coastal storms, Nor'Easters, heavy rains, tropical storms, and hurricanes.

Floods can happen almost anywhere in New Jersey, although they do tend to occur in and around areas near existing bodies of water, such as rivers, streams, and the Atlantic Ocean. The most damaging floods (particularly riverine floods) in New Jersey appear to occur in the northern half of the State. This is a function of several physiographic and physical features of the landscape. Greater geographic relief in the northern half results in flowing water moving down steeper gradients and being naturally or artificially channelized through valleys and gullies.

Development patterns have resulted in denser development in northern New Jersey. In addition, proximity to New York City boosts property values and therefore increases damage dollar totals. Extensive development also leaves fewer natural surfaces available to absorb rainwater, forcing water directly into streams and rivers, swelling them more than when more natural surface buffered the runoff rate. Since the Delaware, Raritan and Passaic Rivers drain more than 90% of the northern New Jersey counties, these rivers and their tributaries are common locations for flooding. The one-percent annual chance of flood hazard zones (both A and V-zones) throughout New Jersey are identified in Figure 5.6-4.






Source:NJDEP 2013FEMAFederal Emergency Management Agency



## Ice Jams

Ice jams are common in the northeast United States and New Jersey is not an exception. In fact, according to the United States Army Corps of Engineers, New Jersey had over 100 ice jam incidents documented between 1780 and 2012 (CRREL 2013). The rivers that experienced the greatest number of ice jams during this time period included the Delaware River (31 reported ice jams) and the South Branch Raritan River (20 reported ice jams).

Figure 5.6-5 presents the number of ice jam incidents in New Jersey during this time period.







## Tsunami

According to a document titled U.S. States and Territories National Tsunami Hazard Assessment: Historical Record and Sources for Waves, the United States Atlantic coast and the Gulf Coast states have experienced



very few tsunamis in the last 200 years. Louisiana, Mississippi, Alabama, the Florida Gulf Coast, Georgia, Virginia, North Carolina, Pennsylvania, and Delaware have no known historical tsunami records. Only six tsunamis have been recorded in the other Gulf and East Coast states. Three of these tsunamis were generated in the Caribbean. Two of these tsunamis were related to a magnitude 7+ earthquake along the Atlantic coast. The other reported tsunami occurred in the Mid-Atlantic States that may have been related to an underwater explosion or landslide (Dunbar and Weaver 2008).

Unlike the Atlantic and Gulf Coasts, the Pacific territories, Puerto Rico, and the United States Virgin Islands have a moderate to very high tsunami hazard. The Pacific territories, including Guam, American Samoa and the Northern Marianas, all experience tsunamis and mostly have a moderate hazard. Studies show that Washington, Oregon, California, Puerto Rico, and the United States Virgin Islands have a high tsunami hazard (Dunbar and Weaver 2008).

Tsunami and tsunami-like waves that have impacted the East Coast were analyzed by Lockridge et al. NOAA's National Geophysical Data Center (NGDC) compiled a listing of all tsunamis and tsunami-like waves of the eastern United States and Canada. Forty-nine potential tsunami events have been identified as possibly impacting the East Coast of the United States between 1668 and 2008. Of these events, eight were categorized as definite or probable tsunamis (NOAA NGDC 2013).

The following present the most significant tsunami threats to the East Coast of the United States:

- *Mid-Atlantic Ridge*—The closest tectonic boundary to the East Coast is the spreading Mid-Atlantic Ridge, which contains numerous faults. However, according to the Maine Geological Survey, tsunamis are more likely to occur at convergent margins. In the Caribbean Sea, there is a convergent plate boundary and a region with a higher probability of generating earthquakes that could produce tsunamis. Tsunamis could potentially travel to New England from the Caribbean, the Mid-Atlantic Ridge, or from the Canary Islands.
- *Caribbean Islands*—The Caribbean is home to some of the most geologically active areas outside of the Pacific Ocean. Similar to the Indonesian Islands, this area has a subduction zone that is located just north of Puerto Rico. The North American plate is being subducted beneath the Caribbean Plate at the Puerto Rico Trench. This area includes other troughs and areas of plate tectonics that have produced numerous earthquakes, sub-marine landslides, volcanic eruptions, and resulting tsunami activity.
- North Carolina/Virginia Continental Shelf—Although the East Coast is much less likely to be affected by a tsunami than the west coast, tsunami threats do exist. Evidence of a large sub-marine landslide off the coasts of Virginia and North Carolina was found and named the Albemarle-Currituck Slide. This event occurred approximately 18,000 years ago when over 33 cubic miles of material slid seaward from the edge of the continental shelf, most likely causing a tsunami.
- *Canary Islands*—The Canary Islands are a volcanic island-arc chain located in the eastern Atlantic Ocean, just west of the Moroccan coastline. La Palma is the western-most and youngest of the Canary Islands and is volcanically active with three large volcanoes. It is also the location of the most active volcano of the Canary Islands, Cumbre Vieja, which most recently erupted in 1949 and again in 1971. Based on a study of past landslide deposits and existing geology of the volcano, some scientists suggest that the west flank of the Cumbre Vieja may experience failure during a future eruption, resulting in a landslide of a block of 15 to 20 kilometers wide and 15 to 25 kilometers long into the Atlantic Ocean. A sudden landslide of this magnitude could create a large tsunami. Although the flank instability of Cumbre Vieja is noted, other scientists disagree with massive failure scenarios for the western flank of the volcano. These scientists think it would happen in smaller, separate events that would not be capable of triggering a mega-tsunami. The International Tsunami Information Center stated the following in regards to the creation of a mega-tsunami by massive flank failure:



"While the active volcano of Cumbre Vieja on Las Palma is expected to erupt again, it will not send a large part of the island into the ocean, though small landslides could occur" (State of Maine 2013).

No mega tsunamis have occurred in the Atlantic or Pacific Oceans in recorded history. The colossal collapses of Krakatau or Santorini generated catastrophic waves in the immediate area but hazardous waves did not propagate to distant shores. Numerical and experimental models on such events and of the Las Palma event verify that the relatively short waves from these small occurrences do not travel as do tsunami waves from a major earthquake (State of Maine 2013).

#### Advisory Base Flood Elevation Maps

Prior to Superstorm Sandy, FEMA had begun a coastal flood study to update FIRMs for portions of New York and New Jersey using improved methods and data to better reflect coastal flood risk. The studies included data that were collected and analyzed over a number of years. After Superstorm Sandy in order to help in rebuilding and recovery efforts, FEMA released Advisory Base Flood Elevation (ABFE) maps which were based on best available data and the partially completed coastal flood study for the Atlantic coastal communities. FEMA is currently in the process of releasing preliminary work maps that include full results of the coastal flood study (FEMA Region II 2014).

ABFEs provide a better picture of current flood risk than the existing FIRMs. The new ABFEs are the recommended elevation of the lowest floor of a building. Some communities may require that the lowest floor be built above the ABFE. ABFEs more accurately reflect the true 1% annual chance flood hazard elevations in a given area. Following large storms, FEMA performs an assessment to determine whether the 1% annual chance flood event, shown on the effective FIRMs, adequately reflects the current flood hazard. In some cases, FEMA determines that ABFEs need to be produced, based on the age of the analysis and the science used to develop the effective FIRMs. ABFEs are provided to communities as a tool to support in the recovery process. ABFEs will aide in making those communities more resilient to future events (FEMA Region II 2014).

According to RiskMAP, the ABFE maps included delineated advisory flood hazard zones (Advisory Zone V, Advisory Zone A, and Advisory Zone X). The maps also included ABFE elevations for 1% and 0.2% annual chance flood elevations and the areas of Limit of Moderate Wave Action (LiMWA). Area and limit of structurally damaging wave action, and preliminary Superstorm Sandy high water marks will also be listed on the ABFE maps, along with coastal barrier resource areas.

The State of New Jersey has adopted emergency amendments to the Flood Hazard Area Control Act Rules (N.J.A.C. 7:13) which incorporate the use of ABFEs to determine flood elevations. Under these amendments, flood elevations are now determined either using the higher of the ABFE, the effective BFE, or the design flood elevation shown on NJDEP flood maps; or site-specific calculations that demonstrate a different flood elevation (NJDEP 2013a). ABFEs and Advisory Flood Hazard Maps take precedence over previous panels and FIS only in construction and development regulations. Where the SFHA and Advisory Flood Hazard Area maps conflict or overlap, the more stringent requirement will be used (NJDEP 2013b).

According to RiskMAP, the ABFE maps were originally available for 10 New Jersey Counties: Atlantic, Bergen, Burlington, Cape May, Essex, Hudson, Middlesex, Monmouth, Ocean, and Union.

#### Extent

In the case of riverine flood hazard, once a river reaches flood stage, the flood extent or severity categories used by the NWS include minor flooding, moderate flooding, and major flooding. Each category has a definition based on property damage and public threat:



- Minor Flooding minimal or no property damage, but possibly some public threat or inconvenience.
- Moderate Flooding some inundation of structures and roads near streams. Some evacuations of people and/or transfer of property to higher elevations are necessary.
- Major Flooding extensive inundation of structures and roads. Significant evacuations of people and/or transfer of property to higher elevations. (NWS 2011)

The severity of a flood depends not only on the amount of water that accumulates in a period of time, but also on the land's ability to manage this water. The size of rivers and streams in an area and infiltration rates are significant factors. When it rains, soil acts as a sponge. When the land is saturated or frozen, infiltration rates decrease and any more water that accumulates must flow as runoff (Harris 2001).

The frequency and severity of flooding are measured using a discharge probability, which is the probability that a certain river discharge (flow) level will be equaled or exceeded in a given year. Flood studies use historical records to determine the probability of occurrence for the different discharge levels. The flood frequency equals 100 divided by the discharge probability. For example, the 100-year discharge has a 1% chance of being equaled or exceeded in any given year. The "annual flood" is the greatest flood event expected to occur in a typical year. These measurements reflect statistical averages only; it is possible for two or more floods with a 100-year or higher recurrence interval to occur in a short time period. The same flood can have different recurrence intervals at different points on a river.

#### Flood

One hundred-year floodplains (or one-percent annual chance floodplain) can be described as a bag of 100 marbles, with 99 clear marbles and one black marble. Every time a marble is pulled out from the bag, and it is the black marble, it represents a 100-year flood event. The marble is then placed back into the bag and shaken up again before another marble is drawn. It is possible that the black marble can be picked one out of two or three times in a row, demonstrating that a "100-year flood event" could occur several times in a row (Interagency Floodplain Management Review Committee 1994).

The 100-year flood, which is the standard used by most federal and state agencies, is used by the NFIP as the standard for floodplain management and to determine the need for flood insurance. A structure located within a SFHA shown on an NFIP map has a 26% chance of suffering flood damage during the term of a 30-year mortgage.

The extent of flooding associated with a 1% annual probability of occurrence (the base flood or 100-year flood) is used as the regulatory boundary by many agencies. Also referred to as the SFHA, this boundary is a convenient tool for assessing vulnerability and risk in flood-prone communities. Many communities have maps that show the extent and likely depth of flooding for the base flood. Corresponding water-surface elevations describe the water elevation resulting from a given discharge level, which is one of the most important factors used in estimating flood damage.

The term "500-year flood" is the flood that has a 0.2% chance of being equaled or exceeded each year. The 500-year flood could occur more than once in a relatively short period of time. Statistically, the 0.2% (500-year) flood has a 6% chance of occurring during a 30-year period of time, the length of many mortgages.

The 500-year floodplain is referred to as Zone X500 for insurance purposes on FIRMs. Base flood elevations or depths are not shown within this zone and insurance purchase is not required in this zone.



## Tsunami

When a major undersea earthquake occurs near the coast at a shallow depth, a destructive tsunami can be generated. This tsunami could impact near-by coasts within minutes and could travel across entire ocean basins causing damage 1,000 miles away. To notify distant coastal areas, internationally-coordinated tsunami warning systems have been established to provide warning to countries regarding regional-to-distant tsunamis. This information is provided to emergency officials, and as appropriate, directly to the public (International Tsunami Information Centre 2008).

NOAA extensively monitors the Pacific Ocean for tsunamis that could impact Hawaii, Alaska, California, Oregon, and Washington. NOAA's Deep-ocean Assessment and Report Tsunamis (DART) program is part of the United States National Tsunami Hazard Mitigation Program and includes seismic networks, tsunami detection buoys and tidal gauges (Maine Geological Survey 2008).

In the Atlantic Ocean, there is no tsunami monitoring program. Although a monitoring program does not exist, the United States Geological Survey (USGS) operates the United States National Seismograph Network, which is part of the Global Seismic Network that monitors seismic activity around the world. These networks detect seismic events that are capable of producing a tsunami. Soon after an earthquake occurs, activity is recorded by seismographs and sent via satellite to the United States National Seismograph Network in Colorado. There, it is analyzed and warnings, if needed, are issued (Maine Geological Survey 2008).

#### **Previous Occurrences and Losses**

Many sources provided flooding information regarding previous occurrences and losses associated with flooding (riverine, inland, and stormwater) events throughout the State of New Jersey. With so many sources reviewed for the purpose of this Hazard Mitigation Plan (HMP), loss and impact information for many events could vary depending on the source. Therefore, the accuracy of monetary figures discussed is based only on the available information identified during research for this HMP.

As previously stated in the 2011 Plan, NOAA's National Climatic Data Center (NCDC) storm events database reported that New Jersey experienced 1,169 flood events between 1950 and 2009. Between January 1, 2010, and December 31, 2012, an additional 413 flood events occurred in New Jersey. Total property damage was estimated at over \$21.88 billion between January 1, 2010 and December 31, 2012. These events included flash floods, coastal flooding, and floods. According to the Hazard Research Lab at the University of South Carolina's Spatial Hazard Events and Losses Database for the United States (SHELDUS), between 1960 and 2012, 413 flood events occurred within New Jersey. The database indicated that flood events and losses totaled over \$23 billion in property damage and over \$800,000 in crop damage. These events included coastal, coastal flooding, thunderstorms, hail, lightning, severe storms, wind, and flooding. SHELDUS indicated that these events resulted in four injuries and no fatalities. However, these numbers may vary due to the database identifying the location of the hazard event in various forms or throughout multiple counties or regions.

The 2011 Plan discussed specific flooding events that occurred in New Jersey through 2009. For this 2014 Plan update, flood events were summarized between January 1, 2010, and December 31, 2012. Table 5.6-2 includes events discussed in the 2011 Plan and events that occurred between 2010 and 2012. With flood documentation for New Jersey being so extensive, not all sources have been identified or researched. Therefore, Table 5.6-2 may not include all events that have occurred throughout the State.



## Table 5.6-2. Flooding Events in New Jersey

Date(s) of Event	Event Type	Counties Affected	Description
April 1984	Flood	N/A	This flooding event in the Passaic River Basin claimed three lives and caused \$335 million in damages. 9,400 people had to evacuate their homes.
January 19 to 26, 1996	Flash Flood	N/A	Flashing flooding led to larger flooding, particularly along the Delaware and Raritan Rivers.
October 19, 1996	Flooding	N/A	Heavy rain caused widespread and severe flooding throughout northern New Jersey, particularly along the Raritan River and its tributaries, as well as the Rahway and Passaic Rivers.
August 20, 1997	Flash Flood	Atlantic	Torrential rain fell across southeast New Jersey as a low pressure system developed over the Delmarva Peninsula and slowly moved northeast across southern New Jersey. Atlantic County bore the brunt of the storm. Storm totals exceeded eight inches from Estell Manor through Galloway Township, and 13.52 inches at the Atlantic City Airport. This storm caused severe flash flooding with several major roadways washing out and bridges collapsing.
September 16, 1999	Flooding associated with Hurricane Floyd	N/A	Hurricane Floyd caused the largest flood on record along the Raritan River. Extensive flooding occurred throughout central and northern New Jersey. Rainfall totals exceeded 12 inches in several locations, with eight to 10-inch totals widespread.
August 12 to 13, 2000	Flooding	Atlantic, Cape May, Monmouth, Morris, Ocean, Sussex	The combination of a weak onshore flow from a nearly stationary low pressure system off the Delmarva Peninsula and the high tides caused by the full moon led to some minor tidal flooding. A nearly unprecedented torrential downpour (approximately a 1,000-year event) remained stationary for about six hours in eastern Sussex County, resulting in considerable flooding in southeastern Sussex and western Morris Counties. The largest rainfall totals exceeded 12 inches.
July 12, 2004	Flash and Poor Drainage Flood	Burlington	Flash flooding occurred during the late afternoon and evening of July 12, as thunderstorms with torrential downpours kept redeveloping along the Interstate 295 corridor in southern Burlington County. This continued for several hours and resulted in widespread storm totals exceeding six inches across most of the Rancocas Creek Basin. A storm total of 13.20 inches was reported in Tabernacle within a 12-hour period and represented a 1,000-year storm. The excessive rain caused record breaking flash flooding along nearly every stream in the Rancocas Basin and led to the failure or damage of 51 dams in Burlington County. Widespread poor drainage flooding also occurred.
September 18, 2004	Flooding associated with remnants of Hurricane Ivan	Morris, Sussex, Warren	The remnants of Hurricane Ivan interacting with a slowly moving cold front caused widespread, heavy rain to fall during the first half of September 18 in Warren, Sussex, and Morris Counties. Storm totals averaged between three and six inches. This, in combination with even heavier rain in eastern Pennsylvania and southeastern New York State, resulted in the worst flooding along the Delaware River since 1955.
March 2005	Flooding	N/A	Following a major rainstorm at the end of March and another between April 1 and April 3, the Delaware River overflowed its banks, flooding an estimated 3,500 homes and forcing the



Date(s) of Event	Event Type	<b>Counties Affected</b>	Description
			evacuation of more than 5,500 people.
July 17, 2005	Flash Flood	Middlesex	Thunderstorms with torrential downpours caused flash flooding in the Manalapan Brook Basin in southeastern Middlesex County.
June 27, 2006	Flooding	N/A	Several days of heavy rain throughout the Delaware River Basin culminated with major flooding along the Delaware River from June 28 to 30.
April 15 to 16, 2007	Flooding associated with a Nor'Easter	N/A	A spring Nor'Easter dropped four to eight inches of rain over most of New Jersey, resulting in major flooding in New Jersey along the Raritan, Passaic, Millstone, Hackensack, and Great Egg Harbor Rivers.
February 10, 2010	Blizzard	Statewide	For the second time within one week a major winter storm affected New Jersey. Blizzard conditions occurred at times across the extreme southern part of the state during the afternoon and early evening of February 10. Snowfall averaged seven to 15 inches across northwest New Jersey, 12 to 20 inches across central New Jersey, and six to 12 inches across the southern third of New Jersey. Ice accretions were less than one tenth of an inch. Two storm-related deaths occurred in Burlington and Middlesex Counties. See Appendix D for detailed information regarding this event.
March 13 to 21 2010	Flooding	Middlesex, Somerset, Morris, Bergen, Passaic	Four days of rain culminated in major flooding in the Passaic and Raritan Basins and flooding throughout New Jersey. Storm totals averaged between 2.5 to six inches, with the highest amounts in the Raritan and Passaic River Basins. It was the worst flooding in the Raritan Basin since April 2007 and the worst flooding in the Passaic Basin since April 1984. Over 1,000 people were evacuated in Morris and Somerset Counties. In Morris County, about 1,300 homes and businesses were damaged. New Jersey Governor Chris Christie declared a state of emergency on March 14. Periods of rain started during the morning of March 12 and fell at its heaviest on March 13. The heaviest rain fell during the morning of March 13 in the southern third of the State, afternoon in the central part of the State and in the afternoon into the evening in the northern third of the State. Periods of lighter rain persisted into March 14 and 15, which slowed the recession of streams and rivers in the area. The flooding cause over \$81 million in property damage.
September 30 to October 1, 2010	Flooding	Bergen, Camden, Gloucester, Hudson, Hunterdon, Morris, Somerset, Sussex, Union	A series of low pressure systems that moved north along a slowly moving cold front brought heavy rain into the western half of New Jersey on September 30 and October 1. Event precipitation totals ranged between three and seven inches. Totals were lighter along the coastal counties. Several streams and rivers flooded across the area and there was also poor drainage flooding. The first round of heavy rain occurred mainly west of New Jersey during the early morning of September 30. The second and heavier round of precipitation moved in during the evening of September 30 and continued into the morning of October 1. The rain ended by the early afternoon of October 1. The flooding cause approximately \$35,000 in property damage.



Date(s) of Event	Event Type	Counties Affected	Description
December 26-27, 2010	Heavy Snow	Statewide	See Appendix D for detailed information regarding this event.
March 7 to 12, 2011	Flooding	Sussex, Morris, Warren	A slow moving, low pressure, cold front brought between 1.5 and four inches of rain across northern New Jersey from the early morning on March 6 into the early morning of March 7. Melting snow contributed to the runoff. The heaviest rain fell during the late afternoon and evening of March 6. Precipitation turned into snow over the higher terrain of northwest New Jersey during the early morning on March 7 and then ended briefly. In eastern Morris County, sections of the Pompton and Passaic Rivers were still above flood stage when another heavy rain event occurred from the early morning on March 10 into the morning on March 11. An additional two to five inches of rain fell and caused major flooding on both rivers. Governor Chris Christie declared a state of emergency before the start of the second round of heavy rain on March 9. Throughout the state, 683 homes were affected by both flooding events and 207 homes suffered at least major damage. About 1,500 people were evacuated and 2,000 residents were affected by the flood waters. The flooding caused over \$11 million in property damage.
April 16 to 17, 2011	Flooding	Burlington, Camden, Cumberland, Gloucester, Morris, Salem	The strong southeast onshore flow on April 16, combined with the high tides associated with the full moon, produced minor to moderate tidal flooding along the New Jersey coast and moderate to severe flooding of the Delaware Bay in Cape May and Cumberland Counties. Tidal flooding departures increased farther up both Delaware and Raritan Bays. In addition, the funneling effect of southeast winds up the Delaware Bay contributed to increasing tidal departures. The high tide at Reedy Point (New Castle County, Delaware) established an all-time record high. One injury was reported from this event. The flooding cause approximately \$2.75 million in property damage.
August 13 to 16, 2011	Flash Flood	Cumberland, Gloucester, Salem	A series of thunderstorms preceding a cold front brought three to seven inches of rain across a wide portion of New Jersey (less along most of the coast) from overnight on August 13 into the day on August 14. In southern Gloucester, eastern Salem and western Cumberland Counties, rainfall amounts reached seven to 11 inches. Scattered thunderstorms occurred on August 15 and into the morning of August 16. This slowed the recession of rivers and streams in the state. The combined event caused severe flash flooding with dam breaks in southwestern New Jersey and flash flooding and flooding across central and northern New Jersey. The flooding caused over \$50 million in property damage. See Appendix D (FEMA Disasters) for additional information regarding this event.
August 27-28, 2011	Hurricane Irene	Statewide	Hurricane Irene moved made its second landfall as a tropical storm near Little Egg Inlet along the southeast New Jersey coast at around 5:35 a.m. on August 28, 2011 Irene brought tropical- storm force winds, destructive storm surge, and record-breaking freshwater inland flooding across northeast New Jersey that resulted in three deaths, thousands of mandatory, and voluntary evacuations along the coast and rivers from surge and freshwater flooding, and widespread power outages that lasted for up to two weeks. The storm surge of three to five feet caused moderate-to-severe tidal flooding along the ocean side and moderate tidal flooding in Delaware Bay and tidal sections of the Delaware River. Major flooding occurred on the



Date(s) of Event	Event Type	<b>Counties Affected</b>	Description
			Raritan, Millstone, Rockaway, and Passaic Rivers.
			Overall, Irene brought an average rainfall total of 7.03 inches with a maximum rainfall total of 9.85 inches in Cranford (Union County). Another source indicated a maximum rainfall total of 11.27 inches in Freehold. A maximum wind gust of 65 mph was reported in Cape May (Cape May County). A maximum storm surge of 4.63 feet was reported in Sandy Hook. Irene caused approximately \$1 billion in damages in New Jersey and seven deaths in the State. See Appendix D (FEMA Disasters) for additional information regarding this event.
September 7-10, 2011	Remnants of Tropical Storm Lee	Burlington, Camden, Cape May, Atlantic, Ocean	Remnants of Tropical Storm Lee brought three to eight inches of rain to many parts of New Jersey. The heavy rain caused flooding, mainly in west and northwest New Jersey. Most of the damage was reported along the Delaware River, where two homes were destroyed, 24 suffered major damage, 249 suffered minor damage, and 28 others were affected. Many roads were closed throughout the State because of flooding. Freshwater surge caused moderate tidal flooding along sections of the Delaware River. The State had approximately \$11.5 million in damage. See Appendix D (FEMA Disasters) for additional information regarding this event.
August 25 to 26, 2012	Flash Flood	Cape May	A series of slow moving thunderstorms caused flash flooding in Cape May County during the evening and overnight on August 25 and into August 26. Doppler Radar storm total estimates reached around five inches. The flooding caused approximately \$150,000 in property damage.
October 26 - November 8, 2012	Superstorm Sandy	Statewide	<ul> <li>Superstorm Sandy was the costliest natural disaster by far in the State of New Jersey. Recordbreaking high tides and wave action combined with sustained winds as high as 60 to 70 mph with wind gusts as high as 80 to 90 mph to batter the State. Statewide, Sandy caused an estimated \$29.4 billion in damage, destroyed or significantly damaged 30,000 homes and businesses, affected 42,000 additional structures, and was responsible directly or indirectly for 38 deaths. A new temporary inlet formed in Mantaloking (Ocean County) where some homes were swept away. About 2.4 million households in the State lost power. It would take two weeks for power to be fully restored to homes and businesses that were inhabitable. Also devastated by the storm was New Jersey's shellfish hatcheries including approximately \$1 million of losses to buildings and equipment, and product losses in excess of \$10,000 at one location alone.</li> <li>Overall, average rainfall totals were 2.78 inches with a maximum rainfall of 10.29 inches at the Cape May (Cape May County) station. Another source indicated a maximum rainfall total of 12.71 inches in Stone Harbor (Cape May County). A maximum wind gust of 78 mph was reported in Robbins Reef. A maximum storm surge of 8.57 feet was reported in Sandy Hook. Superstorm Sandy caused approximately \$30 billion in damages in New Jersey and caused 12 deaths in the State.</li> </ul>

Source: NOAA-NCDC 2013; New Jersey State HMP 2011 N/A Not Available/Not Applicable



## Ice Jams

There have been 107 reported ice jams in New Jersey over the last 233 years (CRREL 2013). According to the United States Army Cold Regions Research and Engineering Laboratory's (CRREL) database, ice jams have historically formed at various points along the Assunpink Creek, Beaver Brook, Cedar Creek, Delaware River, Flat Brook, Forked River, Great Egg Harbor River, Lamington (Black) River, Maurice River, Musconetcong River, Neshanic River, North and South Branch Raritan River, Passaic River, Pequest River, Raritan River, Stony Brook, Walnut Brook, Wanaque River, and West Brook. Locations of historical ice jam events are indicated in Figure 5.6-5.

The 2011 Plan did not discuss ice jam events; however, for this 2014 Plan update, ice jam events that occurred in the State will be further discussed. Table 5.6-3 lists the total number of ice jam events that occurred in each county in New Jersey. Table 5.6-4 lists the ice jam events that have occurred in New Jersey between 1780 and 2012. Information regarding losses associated with these reported ice jams was limited.

County	Total Number of Ice Jams
Atlantic County	1
Bergen County	0
Burlington County	0
Camden County	0
Cape May County	0
Cumberland County	0
Essex County	0
Gloucester County	0
Hudson County	0
Hunterdon County	24
Mercer County	26
Middlesex County	0
Monmouth County	0
Morris County	6
Ocean County	2
Passaic County	4
Salem County	1
Somerset County	12
Sussex County	11
Union County	0
Warren County	20
Total	107

#### Table 5.6-3. Number of Ice Jams Between 1780 and 2012, by County

Source: CRREL 2013



## Table 5.6-4. Ice Jams in New Jersey Between 1780 and 2012

Event Date	River/Location	County	Description/Losses
March 7, 1904	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 11.2 feet, affected by backwater from ice. Bank- full stage eight feet.
March 8, 1904	Delaware River at Trenton	Mercer	Maximum annual gage height of 22.8 feet, affected by backwater from ice.
January 7, 1905	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 12.5 feet, affected by backwater from ice. Bank- full stage eight feet.
January 26, 1907	Delaware River at Trenton	Mercer	Maximum annual gage height of 9.0 feet, affected by backwater from ice.
March 5, 1920	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 11.5 feet, affected by backwater from ice. Bank- full stage eight feet.
January 22, 1924	Musconetcong River at Hackettstown	Warren	Gage height of 3.44 feet, affected by backwater from ice.
January 23, 1924	Beaver Brook at Belvidere	Warren	Gage height of 3.05 feet, affected by backwater from ice. Additional ice-affected gage height of three feet. Bank-full stage four feet.
December 27, 1924	Beaver Brook at Belvidere	Warren	Gage height of 3.03 feet, affected by backwater from ice. Additional ice-affected gage height of 4.09 feet (maximum for year), reported on February 12. Discharge 600 cfs. Also, ice affected gage heights of 3.03 feet, reported on February 24, and 2.96 feet reported on February 27. Bank-full stage four feet.
February 12, 1925	North Branch Raritan River at Raritan	Somerset	Maximum annual gage height of 9.0 feet, affected by backwater from ice.
February 19, 1926	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 9.52 feet, affected by backwater from ice. Bank- full stage eight feet.
January 16, 1927	Beaver Brook at Belvidere	Warren	Maximum gage height of 3.03 feet, affected by backwater from ice. Bank-full stage four feet.
January 20, 1927	Lamington (Black) River at Pottersville	Somerset	Gage height of 2.83 feet, affected by backwater from ice. Bank-full stage five feet.
January 21, 1927	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 8.01 feet, affected by backwater from ice. Bank- full stage eight feet.
January 3, 1928	Beaver Brook at Belvidere	Warren	Gage height of 3.29 feet, affected by backwater from ice. Additional ice-affected gage height of 3.09 feet was reported on January 22. Bank-full stage four feet.
January 25, 1930	Musconetcong River at Hackettstown	Warren	Maximum annual gage height of 3.58 feet, affected by backwater from ice.
January 26, 1930	Delaware River at Trenton	Mercer	Maximum annual gage height of 8.08 feet, affected by backwater from ice.



Event Date	River/Location	County	Description/Losses
January 27, 1930	Beaver Brook at Belvidere	Warren	Maximum annual gage height of 3.10 feet, affected by backwater from ice. Bank- full stage four feet.
December 19, 1932	Beaver Brook at Belvidere	Warren	Gage height of 2.94 feet, affected by backwater from ice. Bank-full stage four feet.
February 13, 1933	Delaware River at Trenton	Mercer	Gage height of 7.90 feet, affected by backwater from ice.
January 4, 1934	Delaware River at Trenton	Mercer	Gage height of 11.83 feet, affected by backwater from ice. Additional ice- affected gage height of 14.2 feet (maximum for year), reported March 5.
January 31, 1934	Beaver Brook at Belvidere	Warren	Gage height of 3.04 feet, affected by backwater from ice. Additional ice-affected gage height of 3.30 feet (maximum for year), reported on March 3. Bank-full stage four feet.
March 3, 1934	Lamington (Black) River at Pottersville	Somerset	Gage height of 3.33 feet, affected by backwater from ice. Additional ice-affected gage height of 3.51 feet, reported on March 4. Bank-full stage five feet.
March 4, 1934	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 10.05 feet, affected by backwater from ice. Daily mean discharge 2,980 cfs. Bank-full stage eight feet.
March 5, 1934	Flat Brook at Flatbrookville	Sussex	Maximum annual gage height of 6.40 feet, affected by backwater from ice. Discharge 700 cfs. Bank-full stage five feet.
January 25, 1935	Delaware River at Trenton	Mercer	Gage height of 7.12 feet, affected by backwater from ice.
December 26, 1935	Delaware River at Trenton	Mercer	Gage height of 6.57 feet, affected by backwater from ice. Additional ice-affected gage height of 16.12 feet, reported on January 3 and ice-affected gage height of 10.20 feet, reported on January 22.
January 3, 1936	Beaver Brook at Belvidere	Warren	Gage height of 3.24 feet, affected by backwater from ice. Additional ice-affected gage height of 3.10 feet, reported on January 21. Also ice-affected gage height of 3.68 feet, reported on January 26. Bank-full stage four feet.
January 3, 1936	North Branch Raritan River at Far Hills	Somerset	Maximum annual gage height of 4.81 feet, affected by backwater from ice.
January 3, 1936	Wanaque River at Monks	Passaic	Gage height of 1.84 feet, affected by backwater from ice. Additional ice-affected gage height of 1.50 feet, reported on February 15.
January 3, 1936	Lamington (Black) River at Pottersville	Somerset	Maximum annual gage height of 4.19 feet, affected by backwater from ice. Discharge 780 cfs. Bank-full stage five feet.
January 3, 1936	Lamington (Black) River at Pottersville	Somerset	Maximum gage height of 4.19 feet caused by an ice jam reported by the USGS.
January 25, 1936	Musconetcong River at Hackettstown	Warren	Gage height of 4.18 feet, affected by backwater from ice.
January 25, 1936	Maurice River at Norma	Salem	Gage height of 4.01 feet, affected by backwater from ice. Bank-full stage 3.5 feet.
February 16, 1936	Cedar Creek at Lanoka Harbor	Ocean	Maximum peak stage of 6.50 feet due to backwater from ice and tide.
January 28, 1938	Beaver Brook at Belvidere	Warren	Gage height of 3.05 feet, affected by backwater from ice. Additional ice-affected gage height of 3.12 feet reported on January 29. Bank-full stage four feet.



Event Date	River/Location	County	Description/Losses
January 27, 1939	Delaware River at Trenton	Mercer	The gage reported water levels of 4.2 feet due to an ice gorge at the gage. Flood stage is 7.5 feet. The gorge was reported through January 28 and resulted in water levels of 4.1 feet on January 29 due to an ice gorge below the gage.
January 30, 1939	South Branch Raritan River at Stanton	Hunterdon	Gage height of 7.32 feet, affected by backwater from ice. Bank-full stage eight feet.
January 30, 1939	Delaware River at Trenton	Mercer	Gage height of 7.40 feet, affected by backwater from ice.
January 15, 1940	Flat Brook at Flatbrookville	Sussex	Gage height of 5.47 feet, affected by backwater from ice. Bank-full stage five feet.
January 15, 1940	South Branch Raritan River at High Bridge	Hunterdon	Gage height of 10 feet, affected by backwater from ice.
January 15, 1940	Lamington (Black) River at Pottersville	Somerset	Gage height of 3.54 feet, affected by backwater from ice. Bank-full stage five feet.
January 15, 1940	South Branch Raritan River at Stanton	Hunterdon	Gage height of 7.91 feet, affected by backwater from ice. Additional ice-affected gage height of eight feet reported on February 11. Bank-full stage eight feet.
January 16, 1940	Delaware River at Trenton	Mercer	Gage height of 8.12 feet, affected by backwater from ice.
February 15, 1940	Wanaque River at Monks	Passaic	Gage height of 1.92 feet, affected by backwater from ice.
March 8, 1941	Pequest River at Huntsville	Sussex	Maximum annual gage height of 3.25 feet, affected by backwater from ice. Bank- full stage four feet.
February 4, 1942	Delaware River at Trenton	Mercer	Gage height of 6.53 feet, affected by backwater from ice.
December 4, 1942	Beaver Brook at Belvidere	Warren	Gage height of 2.97 feet, affected by backwater from ice. Bank-full stage four feet.
December 22, 1942	Lamington (Black) River at Pottersville	Somerset	Gage height of three feet, affected by backwater from ice. Bank-full stage five feet.
January 5, 1943	Pequest River at Huntsville	Sussex	Gage height of 3.32 feet, affected by backwater from ice. Bank-full stage four feet.
February 16, 1943	Delaware River at Trenton	Mercer	Gage height of 6.82 feet, affected by backwater from ice. Additional ice-affected gage height of 7.99 feet, reported on February 20.
January 10, 1944	Beaver Brook at Belvidere	Warren	Gage height of 3.07 feet, affected by backwater from ice. Additional ice-affected gage height of 3.01 feet reported on February 15. Bank-full stage four feet.
February 15, 1944	South Branch Raritan River at High Bridge	Hunterdon	Maximum annual gage height of 10.39 feet, affected by backwater from ice.
January 4, 1945	Beaver Brook at Belvidere	Warren	Gage height of 3.03 feet, affected by backwater from ice. Additional ice-affected gage height of 3.03 feet reported on January 19. Ice-affected gage height of 3.02 feet was reported on January 20. Bank-full stage four feet.
January 12, 1945	Delaware River at Trenton	Mercer	Gage height of 8.24 feet, affected by backwater from ice. Additional ice-affected gage height of 8.72 feet reported on January 17.
January 17, 1945	Delaware River at Trenton	Mercer	Gage height of 8.72 feet, affected by backwater from ice.



Event Date	River/Location	County	Description/Losses
February 22, 1945	Neshanic River at Reaville	Hunterdon	Gage height of 8.42 feet, affected by backwater from ice. Bank-full stage nine feet.
February 22, 1945	South Branch Raritan River at Stanton	Hunterdon	Gage height of 7.73 feet, affected by backwater from ice. Bank-full stage eight feet.
February 27, 1945	Passaic River at Chatham	Morris	Maximum annual gage height of 6.67 feet, affected by backwater from ice.
March 4, 1945	Delaware River at Montague	Sussex	Maximum annual gage height of 17.54 feet, affected by backwater from ice. Additional ice-affected gage height of 15.42 feet was reported on February 28.
December 20, 1945	Delaware River at Trenton	Mercer	Gage height of 8.67 feet, affected by backwater from ice. Additional ice-affected gage height of 11.01 feet (maximum for year), reported on December 26.
December 25, 1945	South Branch Raritan River at High Bridge	Hunterdon	Maximum annual gage height of 9.75 feet, affected by backwater ice.
December 25, 1945	Lamington (Black) River at Pottersville	Somerset	Gage height of 3.66 feet, affected by backwater from ice. Discharge 450 cfs. Bank-full stage five feet.
December 26, 1945	Beaver Brook at Belvidere	Warren	Maximum annual gage height of four feet, affected by backwater from ice. Bank- full stage four feet.
December 26, 1945	Walnut Brook at Flemington	Hunterdon	Gage height of 2.32 feet, affected by backwater from ice. Bank-full stage three feet.
December 26, 1945	Wanaque River at Monks	Passaic	Gage height of 1.87 feet, affected by backwater from ice.
December 26, 1945	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 9.06 feet, affected by backwater from ice. Bank- full stage eight feet.
December 27, 1945	Delaware River at Montague	Sussex	Gage height of 14.7 feet, affected by backwater from ice.
February 10, 1947	Delaware River at Trenton	Mercer	Maximum gage height of 7.9 feet, affected by backwater from ice.
January 25, 1948	Musconetcong River at Bloomsbury	Hunterdon	Gage height of 3.64 feet, affected by backwater from ice. Bank-full stage four feet.
February 19, 1948	South Branch Raritan River at Stanton	Hunterdon	Gage height of 8.54 feet, affected by backwater from ice. Bank-full stage eight feet.
February 20, 1948	North Branch Raritan River at Raritan	Somerset	Maximum annual gage height of 9.39 feet, affected by backwater from ice.
February 21, 1948	Delaware River at Montague	Sussex	Gage height of 17.88 feet, affected by backwater from ice.
February 24, 1948	Passaic River at Chatham	Morris	Maximum annual gage height of 6.65 feet, affected by backwater from ice. Additional ice-affected gage height of 6.3 feet reported on February 20. Estimated daily mean discharge 1,000 cfs.
December 30, 1948	West Brook at Wanaque	Passaic	Gage height of 2.65 feet, affected by backwater from ice. Discharge 388 cfs.
December 21, 1951	Delaware River at Trenton	Mercer	Gage height of 9.48 feet, affected by backwater from ice.
January 21, 1954	South Branch Raritan River at High Bridge	Hunterdon	Maximum annual gage height of 8.97 feet, affected by backwater from ice.
February 7, 1955	Delaware River at Trenton	Mercer	Gage height of 7.27 feet, affected by backwater from ice.

Section 5.6. Flood



Event Date	River/Location	County	Description/Losses
January 23, 1957	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 6.74 feet, affected by backwater from ice. Bank- full stage eight feet.
March 2, 1958	Pequest River at Pequest	Warren	Maximum annual gage height of 3.61 feet, affected by backwater from ice. Bank- full stage four feet.
January 2, 1959	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 7.59 feet, affected by backwater from ice. Discharge 2,310 cfs. Bank-full stage eight feet.
January 6, 1959	Great Egg Harbor River at Folsom	Atlantic	Gage height of 4.72 feet, affected by backwater from ice. Bank-full stage five feet.
January 21, 1959	Lamington (Black) River at Pottersville	Somerset	Maximum annual gage height of 3.64 feet, affected by backwater from ice. Bank- full stage five feet.
January 22, 1959	Pequest River at Pequest	Warren	Gage height of 3.53 feet, affected by backwater from ice. Discharge 640 cfs. Bank-full stage four feet.
January 1, 1961	Neshanic River at Reaville	Hunterdon	Maximum annual gage height of 7.07 feet, affected by backwater from ice. Bank- full stage nine feet.
February 19, 1961	South Branch Raritan River at Stanton	Hunterdon	Maximum annual gage height of 7.28 feet, affected by backwater from ice. Bank- full stage eight feet.
February 20, 1961	Flat Brook at Flatbrookville	Sussex	Maximum annual gage height of 5.67 feet, affected by backwater from ice. Bank- full stage five feet.
February 22, 1961	Passaic River at Chatham	Morris	Maximum annual gage height of 6.59 feet, affected by backwater from ice.
January 15, 1968	Delaware River at Trenton	Mercer	An ice jam was observed at Trenton along the Delaware River.
February 15, 1971	Delaware River at Montague	Sussex	The USGS reported an ice jam on February 15 at Montague on the Delaware River. The estimated water discharge was 10,000 cfs. Maximum gage height was 12.57 feet.
January 18, 1994	Assunpink Creek at Clarksville	Mercer	A flood warning was issued for this USGS gage. The river gage was reading 6.75 feet at 1:40 a.m. and had risen three feet since 7 p.m. due to an ice jam.
January 28, 1994	South Branch Raritan River at High Bridge	Hunterdon	Maximum peak stage of 14.26 feet on January 28 as a result of an ice jam
February 1, 1994	Delaware River at Trenton	Mercer	This jam was approximately one mile long with a backwater of approximately three to four feet. Downstream, the jam was a smooth ice cover about 0.5 to one mile long.



Event Date	River/Location	County	Description/Losses
January 21, 1996	Delaware River at Trenton	Mercer	Ice jams were reported on the Susquehanna, Delaware, and Schuylkill Rivers on January 21. These jams caused severe flooding in Trenton, forcing the evacuation of 3,000 people in the area. Two local people drowned while seven other deaths in the State of Pennsylvania were reported. Ten thousand people in the Wilkes- Barre region were evacuated. The Delaware had risen 12 feet in 10 hours while the Susquehanna crested at 12 feet above flood stage. In Avondale, 109 people were evacuated by boat while another 90 were evacuated from the Bridgeport Towers apartments. Front Street row houses were evacuated as well. This began with a winter storm dumping incredible amounts of snow across Pennsylvania. Of the 40 inches on the ground, 28 inches of it melted. There were also high winds reaching 58 mph.
January 18, 1999	Multiple locations	Sussex	The combination of showers and thunderstorms with heavy rain, already saturated ground, and ice jams along area streams caused flooding and led to the collapse of the foundations of three homes in Hamburg Borough and Andover Township.
January 22, 1999	Delaware River at Depue Island	Warren	An ice jam formed slightly downstream of an existing jam on the Delaware River. Park Rangers reported that it extended from Depue Island north past Tocks Island to Poxono Island. The ice in the Delaware Water Gap was beginning to break up and was predicted to move out later that day.
February 7, 2003	South Branch Raritan River at High Bridge	Hunterdon	A small ice jam formed on the South Branch Raritan River near High Bridge.
February 17, 2003	Forked River at Forked River	Ocean	An ice jam about 300 to 400 yards long formed on the canal leading from Barnegat Bay to the Oyster Creek generating station. The head of the jam was at the Route 1 bridge. The jam in this tidal area was composed of broken ice pieces and slush ice. Its formation occurred after extremely cold air temperatures and a large snowstorm. The jam was restricting primary cooling water flow to the generating plant. Mechanical removal of the jam from the upstream end towards the downstream end was recommended.
February 19, 2003	Delaware River at Trenton	Mercer	The NWS reported an apparent ice jam on the Delaware River at Trenton on February 19. A significant within-banks rise was occurring on the lower main stem of the Delaware River at Trenton, most likely due to an ice jam at the Calhoun Street Bridge. The stage was 15.1 feet at 6 p.m The river had risen over two feet since noon but had stabilized at about 15 feet during the evening.
February 23, 2003	Passaic River at Chatham	Morris	The NWS reported an ice jam along the Passaic River which caused some minor flooding near Chatham. The river stopped rising just above flood stage and stabilized.
February 24, 2003	Passaic River at Chatham	Morris	Maximum gage height of 6.35 feet due to ice effects.
January 31, 2004	Delaware River at Trenton	Mercer	The NWS noted that there was an ice jam north of Trenton on the Delaware River.



Event Date	River/Location	County	Description/Losses	
February 6, 2004	Passaic River at Chatham	Morris	Maximum peak stage of 10.93 feet as a result of an ice jam. The average daily discharge was estimated to be 490 cfs.	
February 6, 2004	Green Brook	Morris	An ice jam developed on the Green Brook. Dynamite was used to break the jam.	
February 6, 2004	Stony Brook at Princeton	Mercer	Maximum gage height of 5.73 feet due to an ice jam. The average daily discharge was estimated to be 280 cfs.	
February 7, 2004	Raritan River at Raritan	Somerset	Maximum peak stage of 11.32 feet as a result of an ice jam. The average daily discharge was estimated to be 3,150 cfs.	
February 14, 2007	Pequest River at Belvidere	Warren	An ice jam formed between two dams on the Pequest River. The lower dam was just above the confluence with the Delaware River, and the upper dam was about 200 yards upstream. Based on descriptions of the ice and local weather, the jam was a freeze-up jam. The ice backed up water into local residents' basements.	
January 27, 2009	Delaware River at Minisink Island	Sussex	An ice jam at Minisink Island was reported to be creating several feet of backwater.	
January 27, 2011	January 27, 2011 Delaware River at Trenton		An ice jam formed downstream from the gaging station at the Trenton Makes Bridge. Water levels increased from nine feet to 13 feet. The ice jam became more restrictive and pushed water up another two feet at the gage.	
January 31, 2011	Delaware River at Montague	Sussex	Solid ice cover was observed upstream from the Milford-Montague toll bridge. There was significant backwater from ice at the gaging station. There was an ice jam upstream in the area of Mashipacong Island.	

Source: CRREL 2013; NOAA-NCDC 2013

Note: Bank-full stage is associated with the flow that just fills the channel to the top of its banks and at a point where the water begins to overflow onto a floodplain

cfs cubic feet per second

mph miles per hour

N/A Not applicable/Not available

NWS National Weather Service

USGS United States Geological Survey



## Tsunami

While the probability of a large tsunami impacting the coast of New Jersey is very small due to the position of New Jersey on the trailing edge of the North Atlantic Plate, the Mid-Atlantic region has been subjected to minor tsunami action over the past 250 years and perhaps significant tsunami action over the last geologic period.

Lockridge, et al. (2002) analyzed tsunami and tsunami-like waves that have impacted the East Coast of the United States NOAA's NGDC compiled a listing of all tsunamis and tsunami-like waves of the eastern United States and Canada. Thirty-nine potential tsunami events have been identified as possibly impacting the East Coast of the United States between 1668 and 2012. Of these events, four are categorized as definite or probable tsunamis.

The NGDC identified six potential tsunami events that have possibly impacted the State of New Jersey. Of those six events, one was categorized as a probably tsunami. Table 5.6-5 describes potential tsunami events that have impacted the State of New Jersey.

## Table 5.6-5. Potential Tsunami Events in New Jersey, 1821-2012

Event Date	Source Location	County	Description/Losses
September 3, 1821	North Carolina	N/A	A hurricane passed over the Outer Banks of North Carolina and over the Delmarva Peninsula. It entered Cape May County where it traveled up the Garden State Parkway. Miles of sandbars were exposed the next morning. A dull roar approached and then a solid mass of wind and rain came tearing great pines from the ground and moving houses from their foundations. A wall of water struck that carried away people and animals.
August 10, 1884	Philadelphia, PA	N/A	A 5.6 earthquake generated a tsunami that was reported from Philadelphia, Trenton, and Highlands. In Trenton, the water in the city reservoir was agitated and a small tidal wave was noticed on the canal and feeder. In Highlands, two men were fishing and felt as if the water was had gone out from under their boat and it was grating on the sand.
September 8, 1889	Asbury Park, NJ	Monmouth	This event occurred during the Mudhen Hurricane. Unusually high waves were reported between September 8 and 10 in the Mid-Atlantic Coast. In New Jersey, these waves were reported in Asbury Park, Atlantic City, Sea Isle City, Coney Island, Long Island, Staten Island and other exposed points.
September 1, 1895	High Bridge, NJ	Hunterdon	A 4.3 earthquake centered near High Bridge was felt over a large area to the northeast and southwest. The earthquake was felt from Maine to Virginia. The earthquake knocked articles from shelves and rocked buildings in several towns in New Jersey, Pennsylvania, and New York. In Asbury Park, NJ, plaster was knocked from walls. The earthquake caused a tsunami-like wave on Long Island. There was one run-up associated with this event. It caused one injury.
June 9, 1913	Longport, NJ	Atlantic	It was reported that heavy tides were associated with this event. There were no reports of storms or earthquakes in the northeast United States on this date. Damage in Longport occurred at the Thoroughfare waterfront when a 250-foot section of the embankment at 23rd Street was carried away. The washout extended to within 15 feet of the near rail line. The tide tore away the wharf at the Schurch chandlery store and it undermined the soil from the building. The Lavine Wharf was completely torn away. This event caused \$10,000 in damage. There was one injury associated with this event.
August 19, 1931	Atlantic City, NJ	Atlantic	There was a sudden and brief onset of 3-meter waves in Atlantic City. Reports state that the surf was rough the day of the event and the waves rolled in shortly before noon. The waves arrived during high tide. There were other high wave events in the region, causing four people to drown. The weather bureau attributed this event to a tropical storm north of Puerto Rico.

Source: Lockridge et al. 2002, NOAA NGDC 2013

N/A Not applicable/Not available

NJ New Jersey

PA Pennsylvania



According to the 2008 NOAA study (U.S. States and Territories National Tsunami Hazard Assessment: Historical Record and Sources for Waves), tsunami events and losses were summarized for the Atlantic Region. Table 5.6-6 is a summary of their findings for the Atlantic Region. Figure 5.6-6 shows the number of tsunami events and total number of events causing run-up heights from 0.1 meters to greater than three meters for the United States and its territories in the Atlantic, Gulf Coast, Puerto Rico, and the United States Virgin Islands.

The table indicates that New Jersey has experienced six tsunami events with any observed run-up. Run-up is a measurement of the height of the water onshore observed above a reference sea level. Tsunami run-up occurs when a peak in the tsunami wave travels from the near-shore region onto shore. There were no reported deaths or injuries associated with these events.


## Table 5.6-6. Summary of Tsunami Events and Losses in the Atlantic Region

Location (and year of first confirmed report)	Tsunami events with observed run-up	Events with undetermined run-up heights	Events with run-ups 0.01 to 0.5 meters	Events with run-ups 0.51 to 1 meters	Events with run-ups 1.01 to 3 meters	Events with run-ups > 3meters	Total number of run-ups for all events	Reported Deaths	Damages (in \$ million)
Maine (1929)	1	1	-	-	-	-	3	-	-
New Hampshire (1929)	1	1	_	-	-	-	1	_	-
Massachusetts (1929)	1	1	—	-	_	-	2	—	-
Rhode Island (1929)	2	1	1	-	-	-	3	-	-
Connecticut (1964)	1	1	-	-	-	-	1	-	-
New York (1895)	2	1	1	-	-	-	7	-	-
New Jersey (1918)	6	3	2	1	-	-	8	-	-
Pennsylvania	-	-	-	-	-	-	-	-	-
Delaware	-	-	-	-	-	-	-	-	-
Maryland (1929)	1	N/A	1	-	-	-	1	-	-
Virginia	-	-	-	-	-	-	-	-	-
North Carolina	-	-	-	-	-	-	-	-	-
South Carolina (1886)	2	1	1	-	-	-	2	-	-
Georgia	-	-	-	-	-	-	-	-	-
Florida (1886)	4	3	1	-	-	-	5	-	-
Atlantic Coast Totals	21	13	7	1	0	0	33	0	\$0

Source: Dunbar and Weaver 2008

*N/A Not applicable/Not available* 

Note: Pennsylvania, Delaware, Virginia, North Carolina and Georgia have not experienced any tsunami events.





*Note:* New Jersey is shown has experiencing six total tsunami events, with three having undetermined run-up heights; two having run-ups of 0.01 to 0.5 meters; and one having run-up of 0.51 to 1 meter

# FEMA Disaster Declarations

Between 1954 and 2012, FEMA declared that the State of New Jersey experienced 26 flood-related disasters (DR) or emergencies (EM) classified as one or a combination of the following disaster types: severe storms, winter storms, snowstorms, coastal storms, flash flooding, heavy rains, tropical storms, hurricanes, high winds, ice jams, wave action, high tide, and tornadoes. Generally, these disasters cover a wide region of the State; therefore, they may have impacted many counties. However, not all counties were included in the disaster declarations as determined by FEMA (FEMA 2013b).

Based on all sources researched, known flooding events that have affected New Jersey and were declared a FEMA disaster, are identified in Table 5.6-7. This table provides information on the FEMA disaster declarations for flooding, including disaster number, disaster type, declaration and incident dates, and counties included in the declaration. Figure 5.6-7 illustrates the number of FEMA-declared disasters by county.

Detailed information pertaining to each of the declared disasters is provided in Appendix D of this Plan.

Source: Dunbar and Weaver 2008

# Table 5.6-7. FEMA Flood-Related Disaster Declarations (1954 to 2012)

			County																						
Disaster Number	Disaster Type	Declaration Date	Incident Period	Atlantic	Bergen	Burlington	Camden	Cape May	Cumberland	Essex	Gloucester	Hudson	Hunterdon	Mercer	Middlesex	Monmouth	Morris	Ocean	Passaic	Salem	Somerset	Sussex	Union	Warren	Number of Counties Impacted
41	Hurricane, Floods	8/20/1955	8/20/1955										N	lot A	vail	able									
124	Severe Storm, High Tides, Flooding	3/9/1962	3/9/1962																						
245	Heavy Rains, Flooding	6/18/1968	6/18/1968		X					Χ					X		Х		X		X		X		7
310	Heavy Rains, Flooding	9/4/1971	9/4/1971	Х	Х	X	Х	X	Х	Х	Х	Х	X	X	X	X	Х	Х	Х	Х	Х	X	X	Х	21
402	Severe Storms, Flooding	8/7/1973	8/7/1973							X					X						x		x		4
519	Severe Storms, High Winds, Flooding	8/21/1976	8/21/1976	X				X								x		x							4
701	Coastal Storms, Flooding	4/12/1984	3/28/1984 to 4/8/1984	X	x			X		x						x	X	x	x						8
973	Coastal Storm, High Tides, Heavy Rain, Flooding	12/18/1992	12/10/1992 to 12/17/1992	X	X			X	X	X		X			x	X		X		X	X		x		12
1145	Severe Storms/Flooding	11/19/1996	10/18/1996 to 10/23/1993									x			x		X				x		x		5
1189	Flooding	9/23/1997	8/20/1997 to 8/21/1997	x																					1
1295	Hurricane Floyd	9/18/1999	9/16/1999 to 9/18/1999		x					X			X	X	x		x		X		x		x		9
1337	Severe Storms,	8/17/2000	8/12/2000 to														Х					X			2



			County																						
Disaster Number	Disaster Type	Declaration Date	Incident Period	Atlantic	Bergen	Burlington	Camden	Cape May	Cumberland	Essex	Gloucester	Hudson	Hunterdon	Mercer	Middlesex	Monmouth	Morris	Ocean	Passaic	Salem	Somerset	Sussex	Union	Warren	Number of Counties Impacted
	Flooding and Mudslides		8/21/2000																						
1530	Severe Storms and Flooding	7/16/2004	7/12/2004 to 7/23/2004			x	x																		2
1563	Tropical Depression Ivan	10/1/2004	9/18/2004 to 10/1/2004										x	x								x		x	4
1588	Severe Storms and Flooding	4/19/2005	4/1/2005 to 4/3/2005		X					X	x		x	x			x		x			x		x	9
1653	Severe Storms and Flooding	7/7/2006	6/23/2006 to 7/10/2006										x	x								x		x	4
1694	Severe Storms and Inland and Coastal Flooding	4/26/2007	4/14/2007 to 4/20/2007	x	X	X	X			X		X		X	x		x		x		x	x	x	x	14
1867	Severe Storms and Flooding Associated with Tropical Depression Ida and a Nor'Easter	12/22/2009	11/11/2009 to 11/15/2009	X				x										X							3
1873	Snowstorm	2/5/2010	12/19/2009 to 12/20/2009	x		x	x		x		x							x		x					7
1889	Severe Winter Storm and Snowstorm	3/23/2010	2/5/2010 to 2/6/2010	x		x	x	x	x		x									x					7
1897	Severe Storms and Flooding	4/2/2010	3/12/2010 to 4/15/2010	X	X	X		X	x	X	X		x	X	x	x	x	x	x		x		x		16




													Co	ount	y										
Disaster Number	Disaster Type	Declaration Date	Incident Period	Atlantic	Bergen	Burlington	Camden	Cape May	Cumberland	Essex	Gloucester	Hudson	Hunterdon	Mercer	Middlesex	Monmouth	Morris	Ocean	Passaic	Salem	Somerset	Sussex	Union	Warren	Number of Counties Impacted
1954	Severe Winter Storm and Snowstorm	2/4/2011	12/26/2010 to 12/27/2010	x	X	X		X	x	X		x		X	x	x	x	x	X		x		x		15
4021	Hurricane Irene	8/31/2011	8/27/2011 to 9/5/2011	x	х	X	x	X	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	X	21
4033	Severe Storms and Flooding	9/15/2011	8/13/2011 to 8/15/2011						x		x									x					3
4039	Remnants of Tropical Storm Lee	10/14/2011	9/28/2011 to 10/6/2011										X	x					x			x		x	5
4086	Hurricane Sandy	10/30/2012	10/26/2012 to 11/8/2012	x	X	x	x	x	x	x	x	x	x	x	x	x	x	x	X	x	x	X	x	x	21

Source: FEMA 2013b

Note: Disaster number is issued by FEMA

FEMA Federal Emergency Management Agency





Source: FEMA 2013 FEMA Federal Emergency Management Agency



## **Probability of Future Occurrences**

For the purposes of this Plan, the frequency of hazard events that are declared disasters is defined by the number of declared disaster events for the State over a specified period of time. The historical record indicates the State has experienced 26 flood-related disaster declaration occurrences from 1954 to 2012 (FEMA 2013b). Refer to Table 5.6-7 for a summary of these disasters. Based on these statistics, the State may experience a flood event of disaster declaration proportions once every two (2.2) years. However, some areas of the State are more flood-prone than others. Therefore the frequency and size of flood events varies based on watershed, riverine reach, and location along each reach.

## **Severity**

#### Flooding

As indicated, the principal factors affecting flood damage are flood depth and velocity. The deeper and faster flood flows become, the more damage they can cause. Shallow flooding with high velocities can cause as much damage as deep flooding with slow velocity. This is especially true when a channel migrates over a broad floodplain, redirecting high velocity flows and transporting debris and sediment. Flood severity can also be evaluated by examining peak discharges.

#### Ice Jams

Heavy snowfall and frigid temperatures, followed by sudden warmer temperatures can increase the risk of flooding from snowmelt and ice jams. When river ice piles up at shallow areas, it blocks the flow of water and may cause flooding of nearby areas (Northeast States Emergency Consortium 2013). Damage tends to be localized and relatively minor. However, depending on the magnitude of the ice jam, major damage and losses can result. Ice jams can damage roads, bridges, buildings, and homes. Impacts from ice jams tend to primarily affect areas along rivers, tributaries, or reservoirs. Typically, when ice jam events occur, flooding occurs within the localized areas upstream of the jam (before it breaks) or downstream from the jam when it suddenly moves or releases.

## Tsunamis

Tsunamis are a threat to life and property to anyone that lives near the ocean. The majority of tsunami events have occurred in the Pacific Ocean Basin. Tsunamis are a threat to life and property to anyone that lives near the ocean. According to NOAA's NGDC, between 2000 B.C. and 2013, 2,483 tsunamis were recorded globally. These events caused over 900,000 fatalities worldwide (NGDC 2013). Of those 2,483 events, 272 occurred along the United States shoreline, causing 941 fatalities, 35 injuries, and costing over \$1 billion in damages.

#### Warning Time

## Flooding

Due to the sequential pattern of meteorological conditions needed to cause serious flooding, it is unusual for a flood to occur without warning. Warning times for floods can be between 24 and 48 hours. Flooding is more likely to occur due to a rainstorm when the soil is already wet and/or streams are already running high from recent previous rains. Pre-existing conditions when a storm begins are called "antecedent conditions".



## Ice Jams

The rates of water level rise during an ice jam event can vary from feet per minute to feet per hour. In some cases, communities have many hours of lead time between the time an ice jam forms and its associated flooding. However, in other cases, the lead time can be as little as one hour (NOAA 2013).

## Tsunamis

The National Tsunami Hazard Mitigation Program was formed in 1995 by Congressional action which directed NOAA to form and lead a federal and state working group. The program is a partnership between NOAA, USGS, FEMA, the National Science Foundation (NSF), and the 28 United States coastal states, territories, and commonwealths.

One of the actions outlined by the National Tsunami Hazard Mitigation Program was the development of a tsunami monitoring system to monitor the ocean's activity and make citizens aware of a possible tsunami approaching land. In response, NOAA developed the DART monitoring buoys. To ensure early detection of tsunamis and to acquire data critical for real-time forecasts, NOAA placed DART stations in regions with a history of destructive tsunamis. NOAA completed the original 6-buoy operational array in 2001 and expanded to a full network of 39 stations in March 2008. The information collected by the DART buoys positioned at strategic locations throughout the ocean plays a critical role in tsunami forecasting.

When a tsunami event occurs, the first information available about the source of the tsunami is based only on the available seismic information for the earthquake event. As the tsunami wave propagates across the ocean and successively reaches the DART stations, these systems report sea level measurement information back to the Tsunami Warning Centers. The centers process the information and produce a new and more refined estimate of the tsunami source. The result is an increasingly accurate forecast of the tsunami that can be used to issue watches, warnings, or evacuations.

## **Secondary Hazards**

## Flooding

The most problematic secondary hazard for flooding is bank erosion and landslides, which in some cases can be more harmful than actual flooding. This is especially true in the upper courses of rivers with steep gradients, where floodwaters may pass quickly and without much damage, but scour the banks, edging properties closer to the floodplain or causing them to fall in. Flooding is also responsible for hazards such as landslides, when high flows on steep slopes with saturated soils cause them to fail. Hazardous materials spills are also a secondary hazard of flooding if storage tanks rupture and spill into streams, rivers, or storm sewers.

## Ice Jams

Ice jams in the United States result in three types of situations:

- No flood threat, but environmental and geomorphological impacts possible;
- Freeze-up jams or freezing of mid-season break-up jams that cause chronic flooding problems for the winter season; or
- Break-up ice jams that cause sudden or flash floods (USACE 2013).

Other impacts from ice jams can include structural damage from intake blockages, ice forces, or scouring under ice. Ice jams can cause bank failure, erosion and scour, and channel shifting. Natural habitats for fish, microbial communities, and riverine margins and estuaries may also be impacted by ice jams (USACE 2013).



## Tsunamis

Aside from the tremendous hydraulic force of the tsunami waves themselves, floating debris carried by a tsunami can endanger human lives and batter inland structures. Ships moored at piers and in harbors often are swamped and sunk or are left battered and stranded high on the shore. Breakwaters and piers collapse, sometimes from scouring actions that sweep away their foundation and sometimes because of the direct wave impact. Railroad yards and oil tanks situated near the waterfront are particularly vulnerable. Oil fires frequently result and can be spread by the waves.

Port facilities, naval facilities, fishing fleets, and public utilities are often the backbone of the economy of the affected areas. These resources generally receive the most severe damage. Until debris can be cleared, wharves and piers rebuilt, utilities restored, and fishing fleets reconstituted, communities may find themselves without fuel, food, and employment. Wherever water transport is a vital means of supply, disruption of coastal systems caused by tsunamis can have far-reaching economic effects.

## **Climate Change Impacts**

Providing projections of future climate change for a specific region is challenging. Shorter term projections are more closely tied to existing trends making longer term projections even more challenging. The further out a prediction reaches the more subject to changing dynamics it becomes.

The New Jersey Climate Adaptation Alliance is a network of policymakers, public and private-sector practitioners, academics, non-governmental organizations (NGO), and business leaders aligned to build climate change preparedness in the state of New Jersey. The Alliance is facilitated by Rutgers University, which provides science and technical support, facilitates the Alliance's operations and advances its recommendations. A document titled *Change in New Jersey: Trends and Projections* was developed to identify recommendations for State and local public policy that will be designed to enhance climate change preparedness and resilience in New Jersey (Rutgers 2013).

Temperatures in the Northeast United States have increased 1.5 degrees Fahrenheit (°F) on average since 1900. Most of this warming has occurred since 1970. The State of New Jersey, for example, has observed an increase in average annual temperatures of 1.2°F between the period of 1971-2000 and the most recent decade of 2001-2010 (ONJSC, 2011). Winter temperatures across the Northeast have seen an increase in average temperature of 4°F since 1970 (Northeast Climate Impacts Assessment [NECIA] 2007). By the 2020s, the average annual temperature in New Jersey is projected to increase by 1.5°F to 3°F above the statewide baseline (1971 to 2000), which was 52.7°F. By 2050, the temperature is projected to increase 3°F to 5°F (Sustainable Jersey Climate Change Adaptation Task Force 2013).

Both northern and southern New Jersey have become wetter over the past century. Northern New Jersey's 1971-2000 precipitation average was over five inches (12%) greater than the average from 1895-1970. Southern New Jersey became two inches (5%) wetter late in the 20th century (Office of New Jersey State Climatologist). Average annual precipitation is projected to increase in the region by 5% by the 2020s and up to 10% by the 2050s. Most of the additional precipitation is expected to come during the winter months (New York City Panel on Climate Change [NYCPCC] 2009). Figure 5.6-8 shows the frequency of heavy precipitation events in the northeastern United States




Figure 5.6-8. Frequency of Heavy Precipitation Events in the Northeastern United States, 1910 to 2010

Source: Rutgers Climate Institute 2013

With this increase in frequency of precipitation, New Jersey may experience more flooding events. More intense, frequent flooding could lead to significant habitat loss for wildlife. Salt marshes and estuaries that serve as critical feeding grounds for birds and waterfowl, and as nursery habitats for commercial fish, could be lost (State of New Jersey 2010). Future climate change may also lead to sea level rise which could lead to more frequent and extensive flooding. See Section 5.2 (Coastal Erosion) for detailed information regarding sea level rise (NJDEP 2013c).



# 5.6.2 Vulnerability Assessment

The following sections address assessing vulnerability and estimating potential losses by jurisdiction and to state facilities. To assess the State's exposure to the riverine and coastal flood hazard, a spatial analysis was conducted using the most current 1% annual chance flood hazard areas. This data includes preliminary work maps, preliminary DFIRMs, regulatory DFIRMs, preliminary work maps and Quality 3 (Q3) data. To estimate potential losses, the Hazards U.S. Multi-Hazard (HAZUS-MH) flood model was used. Preliminary work map depth grids provided by NJDEP were used for the coastal areas. Where existing depth grids were not available, approximate depth grids were generated. The digital elevation models (DEM) provided by New Jersey Office of Information Technology were used to generate the depth grids.


Table 5.6-8 lists the data that was utilized for purposes of the vulnerability assessment. Figure 5.6-9 displays the spatial distribution of the different data types.

County	Data (Source and Date)
Atlantic	Preliminary Work Map (6/17/2013), DFIRM for Estell Manor and Weymouth (6/17/2013), and Quality 3 for the remainder of the County (8/2/1982)
Bergen	Preliminary Work Map (6/17/2013) and DFIRM (9/30/2005) for the remainder of the County
Burlington	Preliminary DFIRM (11/30/2010) and Advisory for Bass River and Washington Township (12/13/2012)
Camden	DFIRM Update (6/16/2009)
Cape May	Preliminary Work Map (8/26/2013)
Cumberland	Preliminary Work Map (7/8/2013)
Essex	Preliminary Work Map (7/18/2013) and DFIRM (6/4/2007) for the remainder of the County
Gloucester	DFIRM (1/20/2010)
Hudson	Preliminary Work Map (6/17/2013)
Hunterdon	DFIRM (5/2/2012)
Mercer	Preliminary DFIRM (7/13/2013)
Middlesex	Preliminary Work Map (7/2/2013) and DFIRM (7/6/2010) for the remainder of the County
Monmouth	Preliminary Work Map (6/17/2013) and DFIRM (9/29/2009) for the remainder of the County
Morris	Preliminary DFIRM (6/10/2011)
Ocean	Preliminary Work Map (6/17/2013) and DFIRM (9/29/2006) for the remainder of the County
Passaic	DFIRM (9/28/2007)
Salem	Preliminary Work Map (7/8/2013)
Somerset	DFIRMs (9/28/2007)
Sussex	DFIRMs (9/29/2011)
Union	DFIRMs (9/20/2006) and Advisory (2/25/2013)
Warren	DFIRMs (9/29/2011)
Course, MDE	

#### Table 5.6-8. Data Used for the Risk Assessment

Source: NJDEP 2013c

DFIRM Digital Flood Insurance Rate Map



Source: NJDEP 2013c

20 Miles

10

N



There are no defined stormwater, tsunami or ice jam hazard areas available at this time. Therefore, the vulnerability to these hazards is discussed in a qualitative nature below. As tsunami inundation or hazard areas are developed, they may be used to conduct a spatial analysis to identify the most vulnerable residents and structures in the tsunami hazard zone and be used to focus public education and outreach efforts on these communities. Further, tsunami inundation maps will provide information needed to create evacuation maps.

# Assessing Vulnerability by Jurisdiction

Historically, floods have impacted all 21 New Jersey counties. All counties with local hazard mitigation plans identified flood as a hazard of concern, as listed in Table 5.1-2 in Section 5.1 (State Risk Assessment Overview). Of the five local mitigation plans that ranked risk into high/medium/low categories for this hazard, the following New Jersey counties considered the flood hazard as high: Cape May, Essex, Monmouth, and Somerset counties.

New Jersey is located along the East Coast, is the most densely populated state, and one of the most densely developed states. Approximately 35% of New Jersey is located within the 1% annual chance flood zone, also known as the SFHA. Hudson and Cape May Counties have the greatest percentage of area located within the SFHA. Refer to Table 5.6-9 which summarizes the total area (inclusive of land and water) located in the 1% annual chance flood zone. Please note certain sections of Barnegat Bay and Raritan Bay are excluded from the SFHA provided by NJDEP.

	Total Area (land and	A	-Zone		V-Zone		SFHA
County	water)	Area	% of Total	Area	% of Total	Area	% of Total
Atlantic	610.65	275.68	45.1%	95.37	15.6%	371.05	60.76%
Bergen	239.83	93.32	38.9%	0.00	0.0%	93.32	38.91%
Burlington	820.32	257.29	31.4%	11.49	1.4%	268.78	32.77%
Camden	227.57	44.18	19.4%	0.00	0.0%	44.18	19.41%
Cape May	286.13	190.83	66.7%	68.49	23.9%	259.33	90.63%
Cumberland	501.8	221.22	44.1%	37.53	7.5%	258.75	51.56%
Essex	129.72	43.63	33.6%	1.08	0.8%	44.71	34.47%
Gloucester	336.2	84.79	25.2%	0.00	0.0%	84.79	25.22%
Hudson	51.53	38.17	74.1%	0.38	0.7%	38.55	74.82%
Hunterdon	437.32	40.80	9.3%	0.00	0.0%	40.80	9.33%
Mercer	228.8	42.94	18.8%	0.00	0.0%	42.94	18.77%
Middlesex	316.97	86.28	27.2%	8.59	2.7%	94.86	29.93%
Monmouth	485.68	79.86	16.4%	36.14	7.4%	116.00	23.88%
Morris	481.44	112.12	23.3%	0.00	0.0%	112.12	23.29%
Ocean	757.93	269.20	35.5%	134.93	17.8%	404.13	53.32%
Passaic	198.32	43.47	21.9%	0.00	0.0%	43.47	21.92%
Salem	347.12	149.23	43.0%	22.18	6.4%	171.41	49.38%
Somerset	304.88	54.29	17.8%	0.00	0.0%	54.29	17.81%

#### Table 5.6-9. Area Located in the Flood Hazard Boundaries (Square Miles) by County



	Total Area	A	-Zone	1	V-Zone		SFHA
County	(land and water)	Area	% of Total	Area	% of Total	Area	% of Total
Sussex	535.47	77.39	14.5%	0.00	0.0%	77.39	14.45%
Union	105.38	17.29	16.4%	2.27	2.2%	19.56	18.56%
Warren	362.59	42.34	11.7%	0.00	0.0%	42.34	11.68%
Total	7,765.65	2,264.3	29.2%	418.46	5.4%	2,682.8	34.5%

Source: NJDEP 2013c

Note: Total area includes all land and water.

% percent SFHA Special Flood Hazard Area

The impact of riverine and coastal flooding on life, health, and safety is dependent upon several factors including the severity of the event and whether or not adequate warning time is provided to residents. Exposure represents the population living in or near floodplain areas that could be impacted should a flood event occur. Additionally, exposure should not be limited to only those who reside in a defined hazard zone, but everyone who may be affected by the effects of a hazard event. For example, people may be at risk while traveling in flooded areas, or emergency service access is compromised during an event. The degree of that impact will vary and is not strictly measurable. Table 5.6-10 lists the estimated population located within the 1% flood zones by county using the 2010 Census block centroid. The limitations of this analysis are recognized and should only be used as estimates.

		A-Zo	ne	V-Zo	ne	SFH	A
County	Total 2010 Population	Population	% of Total	Population	% of Total	Population	% of Total
Atlantic	274,549	78,346	28.5%	709	0.3%	79,055	28.8%
Bergen	905,116	61,660	6.8%	0	0.0%	61,660	6.8%
Burlington	448,734	21,409	4.8%	24	0.0%	21,433	4.8%
Camden	513,657	19,573	3.8%	0	0.0%	19,573	3.8%
Cape May	97,265	39,010	40.1%	273	0.3%	39,283	40.4%
Cumberland	156,898	6,027	3.8%	144	0.1%	6,171	3.9%
Essex	783,969	30,461	3.9%	0	0.0%	30,461	3.9%
Gloucester	288,288	8,918	3.1%	0	0.0%	8,918	3.1%
Hudson	634,266	76,136	12.0%	0	0.0%	76,136	12.0%
Hunterdon	128,349	3,232	2.5%	0	0.0%	3,232	2.5%
Mercer	366,513	14,926	4.1%	0	0.0%	14,926	4.1%
Middlesex	809,858	39,575	4.9%	756	0.1%	40,331	5.0%
Monmouth	630,380	55,027	8.7%	3,834	0.6%	58,861	9.3%
Morris	492,276	32,785	6.7%	0	0.0%	32,785	6.7%
Ocean	576,567	89,146	15.5%	2,814	0.5%	91,960	15.9%
Passaic	501,226	24,630	4.9%	0	0.0%	24,630	4.9%
Salem	66,083	15,002	22.7%	30	0.0%	15,032	22.7%

## Table 5.6-10. Estimated Population Exposed to the 1% Annual Chance Flood Events



		A-Zo	ne	V-Zo:	ne	SFH	A
County	Total 2010 Population	Population	% of Total	Population	% of Total	Population	% of Total
Somerset	323,444	18,222	5.6%	0	0.0%	18,222	5.6%
Sussex	149,265	4,672	3.1%	0	0.0%	4,672	3.1%
Union	536,499	34,742	6.5%	0	0.0%	34,742	6.5%
Warren	108,692	3,540	3.3%	0	0.0%	3,540	3.3%
Total	8,791,894	677,039	7.7%	8,584	0.1%	685,623	7.8%

Source: United States Census 2010; NJDEP 2013c % percent SFHA Special Flood Hazard Area

Of the exposed population, the most vulnerable include the economically disadvantaged and those over the age of 65. Economically disadvantaged populations are more vulnerable because they are likely to evaluate their risk and make evacuation decisions based on the net economic impact to their family. Those over 65 are also more vulnerable because they are more likely to seek or need medical attention which may not be available during a flood event, and they may have more difficulty evacuating. Refer to Section 4 (State Profile) which summarizes these vulnerable populations by county.

In terms of the population's vulnerability to flooding, Cape May County has the highest percent of total population located within the SFHA (40%). The following counties have greater than 10% of their population in the SFHA (in descending order): Cape May, Atlantic, Salem, Ocean, Hudson, and Monmouth.

The populations most vulnerable to the tsunami hazard are the elderly, disabled, and very young who reside near beaches, low-lying coastal areas, tidal flats, and river deltas that empty into ocean-going waters. In the event of a local tsunami generated in or near the State, there would be little warning time, so more of the population would be vulnerable. The degree of vulnerability of the population exposed to the tsunami hazard event is based on a number of factors:

- Whether there is a warning system in place;
- How much lead time a warning provides;
- The method for disseminating the warning; and
- Whether the people warned will evacuate.

Damages to buildings can impact a community's economy and tax base. To provide a general estimate of the structural/content replacement value exposure, the 1% annual chance flood boundary was overlaid on HAZUS-MH's default general building stock inventory at the Census Block level for each county. Where the 2010 Census block centroid was located within the flood boundary, the building stock values in that Census block were totaled. There are limitations to this analysis. This methodology was conducted for all counties with the exception of Somerset county. Somerset county's 2013 HMP update included a countywide building update using detailed structure-specific assessor data. The default building stock in HAZUS-MH was replaced with this custom inventory, and the results from their 2013 HMP update provide more accurate estimates. When statewide buildings data (footprints or otherwise) are available, this analysis will be further refined and updated for all counties. Refer to Table 5.6-11 for a summary of this analysis by county.

AT THE STATE OF THE STATE	State of New Jerse	y 2014 Hazard Mi	tigation Pla	n			
Table 5	5.6-11. Estimated	General Building	Stock Expo	sure to the 1%	Annual Cha	nce Flood Event	
		A-Zon	ρ	V-Zo	ne	SFHA	
County	Total Value	Value	% of Total	Value	% of Total	Value	% of Total
Atlantic	\$38,043,171,000	\$13,568,591,000	35.7%	\$369,144,000	1.0%	\$13,937,735,000	36.6%
Bergen	\$154,077,482,000	\$14,201,789,000	9.2%	\$0	0.0%	\$14,201,789,000	9.2%
Burlington	\$62,700,794,000	\$3,477,187,000	5.5%	\$2,523,000	0.0%	\$3,479,710,000	5.5%
Camden	\$70,467,051,000	\$3,169,675,000	4.5%	\$0	0.0%	\$3,169,675,000	4.5%
Cape May	\$24,665,528,000	\$14,843,938,000	60.2%	\$371,706,000	1.5%	\$15,215,644,000	61.7%
Cumberland	\$18,128,613,000	\$1,035,041,000	5.7%	\$31,747,000	0.2%	\$1,066,788,000	5.9%
Essex	\$113,124,687,000	\$9,379,381,000	8.3%	\$0	0.0%	\$9,379,381,000	8.3%
Gloucester	\$33,534,660,000	\$1,446,390,000	4.3%	\$0	0.0%	\$1,446,390,000	4.3%
Hudson	\$82,290,184,000	\$16,008,267,000	19.5%	\$0	0.0%	\$16,008,267,000	19.5%
Hunterdon	\$21,720,513,000	\$839,901,000	3.9%	\$0	0.0%	\$839,901,000	3.9%
Mercer	\$56,194,660,000	\$2,710,602,000	4.8%	\$0	0.0%	\$2,710,602,000	4.8%
Middlesex	\$119,947,782,000	\$7,405,351,000	6.2%	\$61,060,000	0.1%	\$7,466,411,000	6.2%
Monmouth	\$96,235,266,000	\$8,426,671,000	8.8%	\$1,338,961,000	1.4%	\$9,765,632,000	10.1%
Morris	\$86,634,810,000	\$7,092,963,000	8.2%	\$0	0.0%	\$7,092,963,000	8.2%
Ocean	\$73,559,915,000	\$20,304,337,000	27.6%	\$1,377,414,000	1.9%	\$21,681,751,000	29.5%
Passaic	\$66,705,864,000	\$5,951,353,000	8.9%	\$0	0.0%	\$5,951,353,000	8.9%
Salem	\$8,092,037,000	\$1,711,505,000	21.2%	\$18,602,000	0.2%	\$1,730,107,000	21.4%
Somerset	\$83,463,372,709	\$1,839,554,941	2.2%	\$0	0.0%	\$1,839,554,941	2.2%
Sussex	\$20,979,595,000	\$809,907,000	3.9%	\$0	0.0%	\$809,907,000	3.9%
Union	\$79,329,736,000	\$5,978,433,000	7.5%	\$451,275,000	0.6%	\$6,429,708,000	8.1%

\$1,324,338,475,709 Source: HAZUS-MH 2.1; NJDEP 2013c

\$14,442,755,000

\$863,936,000

\$141,064,772,941

Note: The total building replacement cost values (RCV) are for all occupancy types (residential, commercial, industrial, religious, government, and education) and represent both structure and contents.

\$0

\$4,022,432,000

0.0%

0.3%

\$863,936,000

\$145,087,204,941

6.0%

10.7%

% percent

Warren

Total

SFHA Special Flood Hazard Area

The impact of the waves and the scouring associated with debris that may be carried in the water could be very damaging to structures located in the tsunami's path. Structures that would be most vulnerable are those located in the front line of tsunami impact and those that are structurally unsound.

The NFIP data are a useful tool to determine areas vulnerable to flood and severe storm hazards. Table 5.6-12 summarizes the NFIP policies, claims, RL, and SRL properties in each county in 2013 (post-Tropical Storm Irene in 2011, and post-Superstorm Sandy in 2012). Appendix U summarizes this data at the community level. Passaic County has the highest number of SRL properties in the State. Cape May county has the highest number of RL properties in the State.

6.0%

11.0%



County	Number of Policies	Number of Policies V-Zone	Number of Policies A-Zone	Number of Claims	Total Loss Payment	Number RL Properties	Number of SRL Properties
Atlantic	32,382	292	29,676	20,309	\$430,537,978	1,022	89
Bergen	15,694	0	11,866	12,614	\$331,729,130	1,871	178
Burlington	4,201	0	2,411	1,711	\$21,751,013	166	9
Camden	2,486	0	1,370	1,001	\$5,206,049	88	3
Cape May	55,703	573	52,916	26,803	\$361,368,385	2,302	249
Cumberland	835	0	568	971	\$14,158,744	101	2
Essex	4,925	0	3,192	4,627	\$106,961,735	499	73
Gloucester	1,530	0	899	561	\$3,479,499	58	1
Hudson	18,207	0	15,592	4,067	\$132,153,292	415	23
Hunterdon	1,201	0	612	1,301	\$24,582,853	229	33
Mercer	2,500	0	1,327	2,201	\$36,722,471	295	15
Middlesex	4,922	2	2,079	4,142	\$106,803,702	635	83
Monmouth	23,417	285	11,974	19,392	\$800,912,227	1,604	122
Morris	4,841	0	3,306	8,913	\$190,580,089	1,064	275
Ocean	55,011	1,117	48,287	52,007	\$2,182,926,444	1,817	91
Passaic	5,038	0	3,897	13,502	\$277,887,430	1,759	610
Salem	2,259	0	1,921	707	\$6,250,411	47	5
Somerset	3,315	0	1,810	5,275	\$158,616,323	1,031	153
Sussex	432	0	110	182	\$1,687,288	16	0
Union	6,160	0	3,979	5,555	\$95,414,904	728	50
Warren	747	0	406	1,224	\$31,706,999	270	33
Total	245,806	2,269	198,198	187,065	\$5,321,436,966	16,017	2,097

# Table 5.6-12. Current Status of NFIP Policies, Claims, and Repetitive Loss Statistics (2013)

Note: Number of policies and claims are as of October 23, 2013. Claims represent all statuses: Open, Closed with Payment, Closed without payment through October 23, 2013.

Number of repetitive loss properties includes the number of severe repetitive loss properties through June 30, 2013.

NFIP National Flood Insurance Program

RL Repetitive loss property

SRL Severe repetitive loss property

Of the 16,017 RL properties across the State, 11,730 (73%) are single family residences. The remaining properties are classified as the following: two to four family residence (2,229); condominium (238); other residential (417) and non-residential (1,403). Of the 2,097 SRL properties across the State, 1,777 (85%) are single family residences while 279 are two to four family residences and the remaining 41 properties are classified as other residential. This data is current as of June 2013.



A comparison analysis was conducted to understand the changes in overall NFIP statistics from the 2011 Plan to the 2014 Plan update. During this time, New Jersey experienced two major flood disaster declarations: one mainly inland flooding event (Tropical Storm Irene) and one coastal flooding event (Superstorm Sandy). Table 5.6-13 summarizes these findings by county. Ocean County had the highest increase in the number of policies after these events. Passaic County had the highest increase in the number of SRL properties (161), followed closely by Bergen (136), Morris (110), Monmouth (107) and Somerset (91). Essex county is the only county with a decrease in the number of SRL properties. Refer to Table 5.6-13 which summarizes this data and indicates the change over the years for each county.

A comparison analysis was also conducted to understand the changes in repetitive loss and severe repetitive loss properties across the State pre- and post-Tropical Storm Irene and pre- and post-Superstorm Sandy. Tables 5.6-14 and 5.6-15 summarize these findings by county and Figures 5.6-10 to 5.6-15 illustrate these findings.

The comparison between pre- and post-Tropical Storm Irene statistics indicates that Passaic County had the highest increase in the number of SRL properties (211), followed closely by Morris (141), Bergen (104) and Somerset (90) Counties. Essex county is the only county with a decrease in the number of SRL properties from April 2011 to August 2012. All counties experienced an increase in the number of policies. As shown in Table 5.6-14, Monmouth, Cape May, Bergen, Morris, and Hudson counties experienced the greatest increase.

The comparison between pre-and post-Superstorm Sandy statistics indicates that Monmouth county had the highest increase in the number of SRL properties (85), followed by Cape May (57), Ocean (38), Atlantic (37), and Bergen (32) Counties. Ocean, Monmouth, Hudson, and Atlantic counties experienced an increase in greater than 1,000 policies per county, with Ocean and Monmouth exceeding 2,000. Meanwhile Cumberland, Somerset and Warren Counties saw a decrease in the number of NFIP policies. Ocean, Atlantic, Monmouth and Cape May experienced the highest number of claims from pre- to post-Superstorm Sandy.

## Table 5.6-13. Comparison of NFIP Statistics from 2011 to 2013

	Ň	umber of Policie	S	j	Number of Clai	ms	Numb	oer of Repetitiv Properties	e Loss		of Severe R oss Properti	
County	2011	2013	Change	2011	2013	Change	2011	2013	Change	2011	2013	Change
Atlantic	31,749	32,382	633	8,835	20,309	11,474	884	1,022	138	47	89	42
Bergen	13,885	15,694	1,809	6,983	12,614	5,631	817	1,871	1,054	42	178	136
Burlington	4,281	4,201	-80	1,221	1,711	490	103	166	63	2	9	7
Camden	2,409	2,486	77	789	1,001	212	67	88	21	2	3	1
Cape May	53,986	55,703	1,717	16,524	26,803	10,279	2,072	2,302	230	191	249	58
Cumberland	1,055	835	-220	701	971	270	77	101	24	1	2	1
Essex	3,960	4,925	965	3,153	4,627	1,474	331	499	168	237	73	-164
Gloucester	2,132	1,530	-602	379	561	182	23	58	35	1	1	0
Hudson	16,066	18,207	2,141	1,265	4,067	2,802	102	415	313	5	23	18
Hunterdon	1,423	1,201	-222	997	1,301	304	190	229	39	15	33	18
Mercer	880	2,500	1,620	1,764	2,201	437	259	295	36	4	15	11
Middlesex	4,519	4,922	403	2,295	4,142	1,847	297	635	338	29	83	54
Monmouth	23,781	23,417	-364	7,553	19,392	11,839	680	1,604	924	15	122	107
Morris	4,092	4,841	749	6,510	8,913	2,403	730	1,064	334	165	275	110
Ocean	46,595	55,011	8,416	13,317	52,007	38,690	847	1,817	970	42	91	49
Passaic	4,377	5,038	661	10,749	13,502	2,753	1,359	1,759	400	449	610	161
Salem	2,215	2,259	44	452	707	255	19	47	28	1	5	4
Somerset	2,955	3,315	360	3,861	5,275	1,414	837	1,031	194	62	153	91
Sussex	339	432	93	119	182	63	8	16	8	0	0	0
Union	4,759	6,160	1,401	3,721	5,555	1,834	487	728	241	8	50	42
Warren	826	747	-79	1,054	1,224	170	233	270	37	31	33	2
Total	226,284	245,806	19,522	92,242	187,065	94,823	10,422	16,017	5,595	1,349	2,097	748

Sources: NJ State Hazard Mitigation Plan 2011; NJDEP 2013d

Note: The 2011 policies and claims were current as of June 22, 2011; the repetitive loss and severe repetitive loss properties were current as of April 30, 2011. The 2011 number of repetitive loss properties includes the number of severe repetitive loss properties.

State of New Jersey 2014 Hazard Mitigation Plan



The 2013 claims are all Open, Closed with Payment, or Closed without Payment through October 23, 2013. The repetitive loss and severe repetitive loss properties are current as of June 30, 2013. The 2013 number of repetitive loss properties includes the number of severe repetitive loss properties. NFIP National Flood Insurance Program

Section 5.6. Flood

# Table 5.6-14. Pre- and Post-Tropical Storm Irene NFIP Statistics

	Nun	nber of Poli	cies	Nu	mber of Clai	ms	Numbe	r of Repetiti Properties		Number of Severe Repetitive Loss Properties			
County	Pre- Irene	Post- Irene	Change	Pre- Irene	Post- Irene	Change	Pre- Irene	Post- Irene	Change	Pre- Irene	Post- Irene	Change	
Atlantic	30,895	31,080	185	8,835	9,035	200	884	900	16	47	52	5	
Bergen	14,069	14,752	683	6,983	10,399	3,416	817	1,347	530	42	146	104	
Burlington	3,925	4,126	201	1,221	1,592	371	103	156	53	2	8	6	
Camden	2,310	2,430	120	789	918	129	67	81	14	2	3	1	
Cape May	54,379	55,128	749	16,524	16,761	237	2,072	2,102	30	191	192	1	
Cumberland	792	839	47	701	761	60	77	84	7	1	2	1	
Essex	4,356	4,617	261	3,153	4,508	1,355	331	474	143	237	77	-160	
Gloucester	1,434	1,513	79	379	519	140	23	48	25	1	1	0	
Hudson	16,999	17,538	539	1,265	1,861	596	102	162	60	5	10	5	
Hunterdon	1,088	1,171	83	997	1,292	295	190	229	39	15	34	19	
Mercer	2,333	2,437	104	1,764	2,170	406	259	294	35	4	16	12	
Middlesex	4,002	4,420	418	2,295	3,393	1,098	297	482	185	29	64	35	
Monmouth	20,396	21,226	830	7,553	9,329	1,776	680	820	140	15	37	22	
Morris	4,223	4,762	539	6,510	8,862	2,352	730	1,062	332	165	306	141	
Ocean	52,107	52,510	403	13,317	14,496	1,179	847	904	57	42	53	11	
Passaic	4,494	4,888	394	10,749	13,445	2,696	1,359	1,755	396	449	660	211	
Salem	2,175	2,242	67	452	552	100	19	25	6	1	3	2	
Somerset	3,134	3,315	181	3,861	5,231	1,370	837	1,026	189	62	152	90	
Sussex	326	419	93	119	179	60	8	16	8	0	0	0	
Union	5,523	5,897	374	3,721	5,380	1,659	487	685	198	8	34	26	
Warren	668	758	90	1,054	1,218	164	233	270	37	31	47	16	
Total	229,628	236,068	6,440	92,242	111,901	19,659	10,422	12,922	2,500	1,349	1,897	548	

Source: NJDEP 2013d

# State of New Jersey 2014 Hazard Mitigation Plan



Notes: Pre-Tropical Storm Irene policies and claims are current as of April 30, 2011.

Pre-Tropical Storm Irene severe repetitive loss properties are current as of June 22, 2011.

Post-Tropical Storm Irene policies, claims, repetitive loss and severe repetitive loss properties are current as of August 31, 2012.

The number of repetitive loss properties includes the number of severe repetitive loss properties.

NFIP National Flood Insurance Program


## Table 5.6-15. Pre- and Post-Superstorm Sandy NFIP Statistics

	Nur	nber of Poli	cies	Nu	mber of Clai	ims	Numbe	r of Repetiti Properties	ve Loss	Number of Severe Repetitive Loss Properties			
County	Pre- Sandy	Post- Sandy	Change	Pre- Sandy	Post- Sandy	Change	Pre- Sandy	Post- Sandy	Change	Pre- Sandy	Post- Sandy	Change	
Atlantic	31,080	32,151	1,071	9,035	20,282	11,247	900	1,022	122	52	89	37	
Bergen	14,752	15,558	806	10,399	12,605	2,206	1,347	1,871	524	146	178	32	
Burlington	4,126	4,194	68	1,592	1,706	114	156	166	10	8	9	1	
Camden	2,430	2,480	50	918	967	49	81	88	7	3	3	0	
Cape May	55,128	55,571	443	16,761	26,762	10,001	2,102	2,302	200	192	249	57	
Cumberland	839	835	-4	761	971	210	84	101	17	2	2	0	
Essex	4,617	4,865	248	4,508	4,620	112	474	499	25	77	73	-4	
Gloucester	1,513	1,518	5	519	552	33	48	58	10	1	1	0	
Hudson	17,538	18,883	1,345	1,861	4,062	2,201	162	415	253	10	23	13	
Hunterdon	1,171	1,199	28	1,292	1,301	9	229	229	0	34	33	-1	
Mercer	2,437	2,476	39	2,170	2,197	27	294	295	1	16	15	-1	
Middlesex	4,420	4,853	433	3,393	4,140	747	482	635	153	64	83	19	
Monmouth	21,226	23,232	2,006	9,329	19,378	10,049	820	1,604	784	37	122	85	
Morris	4,762	4,833	71	8,862	8,910	48	1,062	1,064	2	306	275	-31	
Ocean	52,510	54,929	2,419	14,496	51,961	37,465	904	1,817	913	53	91	38	
Passaic	4,888	5,013	125	13,445	13,486	41	1,755	1,759	4	660	610	-50	
Salem	2,242	2,248	6	552	697	145	25	47	22	3	5	2	
Somerset	3,315	3,305	-10	5,231	5,273	42	1,026	1,031	5	152	153	1	
Sussex	419	432	13	179	182	3	16	16	0	0	0	0	
Union	5,897	6,112	215	5,380	5,554	174	685	728	43	34	50	16	
Warren	758	741	-17	1,218	1,224	6	270	270	0	47	33	-14	
Total	236,068	245,428	9,360	111,901	186,830	74,929	12,922	16,017	3,095	1,897	2,097	200	

Source: NJDEP 2013d




Notes:Pre-Superstorm Sandy policies and claims are current as of August 2012.Post-Superstorm Sandy policies, claims, repetitive loss and severe repetitive loss properties are current as of June 2013.NFIPNational Flood Insurance Program




#### Figure 5.6-10. NFIP Repetitive Loss Properties Pre-Tropical Storm Irene

Source: NJDEP 2013d Note: Data is current as of June 2011 NFIP National Flood Insurance Program




## Figure 5.6-11. NFIP Repetitive Loss Properties Post-Tropical Storm Irene/Pre-Superstorm Sandy

Source:NJDEP 2013dNote:Data is current as of August 2012NFIPNational Flood Insurance Program






Source: NJDEP 2013d Note: Data is current as of June 2013 NFIP National Flood Insurance Program




# Figure 5.6-13. NFIP Severe Repetitive Loss Areas Pre-Tropical Storm Irene

Source: NJDEP 2013d Note: Data is current as of June 2011 NFIP National Flood Insurance Program





Source: NJDEP 2013dNote:Data is current as of August 2012NFIPNational Flood Insurance Program



#### Figure 5.6-15. NFIP Severe Repetitive Loss Areas Post-Superstorm Sandy

Source: NJDEP 2013d Note: Data is current as of June 2013 NFIP National Flood Insurance Program



As New Jersey continues to develop, the State may remain vulnerable to the impacts of flood hazards, but with mitigating factors. Much of the undeveloped property within these flood prone hazard areas will likely remain undeveloped, and the State's priority to decrease the number of RL and SRL properties with post-disaster funding will contribute to decreasing vulnerable structures in the future.

The release of FEMA's advisory Flood Hazard Areas and preliminary work maps in addition to the changes to New Jersey's Flood Hazard Area Control Act will help mitigate the impacts of these events. The revisions to the Act set new minimum elevation standards for the reconstruction of houses and buildings in areas that are vulnerable to flooding. Through continued public outreach and education, people are becoming increasingly aware that measures, such as elevating their homes or using innovative stormwater management techniques like the installation of rain gardens, will help mitigate the impacts of flood hazards.

Construction and reconstruction within flood hazard areas will be influenced as a result of BW-12. Property owners are being strongly encouraged by NJDEP to consider long-term insurance costs when undertaking reconstruction or elevation of damaged buildings. A relatively small investment to raise the lowest floor of a building an additional foot or two may translate into significant future flood insurance savings. This will positively contribute to decreasing the vulnerability of structures in the flood hazard areas.

# **Assessing Vulnerability to State Facilities**

To assess the vulnerability of the state-owned and -leased facilities provided by New Jersey's Office of Management and Budget (NJ OMB), an analysis was conducted using the 1% annual chance flood hazard areas. Using geographic information system (GIS) software, these hazard areas were overlaid with the state facility data to determine the number of state facilities vulnerable. Table 5.6-16 summarizes the state-owned and -leased facilities vulnerable by county, and Table 5.6-17 summarizes the facilities by state agency.

Overall, there are 157 state-owned or -leased buildings (5.8%) that are vulnerable to the 1% flood hazard (A and V-zones). The greatest numbers of vulnerable State buildings are located in Essex and Ocean counties. There are 10 state-owned buildings located in the V-zone but no state-leased buildings located in the V-zone. The NJDEP has the greatest number of buildings vulnerable to the flood hazard. The NJDEP operates numerous flood control and water assets which accounts for the large number of structures in the flood zone. Refer to Tables 5.6-16 and 5.6-17 below which summarize these findings by county and state agency, respectively.

There are 1,707 critical facilities and infrastructure located in the 1% flood hazard area (A and V-zones). Of these, 957 are dams. Excluding dams from the analysis, which by default are located in flood hazard areas, Hudson county has the greatest number of vulnerable critical facilities and infrastructure. Of all the facility types, schools have the greatest number of vulnerable structures (total of 154), followed by emergency medical services (EMS) (125) and fire (122). Table 5.6-18 summarizes the number of critical facilities and infrastructure located in the hazard area by facility type.

## Table 5.6-16. State Building Exposure to the 1% Annual Chance Flood Hazard, by County

		A-Z	one			V-Zon	e		SFHA							
	(	Owned		Leased	0	wned	Lea	sed	(	Owned		Leased		Total		
County	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value		
Atlantic	6	\$10,527,434	6	\$66,132,108	1	\$5,887,331	0	0	7	\$16,414,766	6	\$66,132,108	13	\$82,546,873		
Bergen	1	\$1,018,773	0	\$0	0	\$0	0	0	1	\$1,018,773	0	\$0	1	\$1,018,773		
Burlington	10	\$7,393,114	0	\$0	0	\$0	0	0	10	\$7,393,114	0	\$0	10	\$7,393,114		
Camden	7	\$2,277,759	1	\$9,319,764	0	\$0	0	0	7	\$2,277,759	1	\$9,319,764	8	\$11,597,523		
Cape May	5	\$1,352,135	1	\$822,003	0	\$0	0	0	5	\$1,352,135	1	\$822,003	6	\$2,174,139		
Cumberland	19	\$1,473,484	0	\$0	1	\$394,669	0	0	20	\$1,868,153	0	\$0	20	\$1,868,153		
Essex	30	\$180,520,018	2	\$12,948,409	0	0	0	0	30	\$180,520,018	2	\$12,948,409	32	\$193,468,426		
Gloucester	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0		
Hudson	9	\$8,430,357	1	\$21,504,792	1	\$29,383,306	0	0	10	\$37,813,663	1	\$21,504,792	11	\$59,318,455		
Hunterdon	3	\$828,444	0	\$0	0	\$0	0	0	3	\$828,444	0	\$0	3	\$828,444		
Mercer	2	\$53,488,685	3	\$42,717,879	0	\$0	0	0	2	\$53,488,685	3	\$42,717,879	5	\$96,206,564		
Middlesex	0	\$0	1	\$451,732	0	\$0	0	0	0	\$0	1	\$451,732	1	\$451,732		
Monmouth	0	\$0	0	\$0	1	\$110,222	0	0	1	\$110,222	0	\$0	1	\$110,222		
Morris	1	\$12,401	0	\$0	0	\$0	0	0	1	\$12,401	0	\$0	1	\$12,401		
Ocean	28	\$10,558,275	1	\$580,145	6	\$1,761,534	0	0	34	\$12,319,809	1	\$580,145	35	\$12,899,954		
Passaic	0	\$0	1	\$8,015,914	0	\$0	0	0	0	\$0	1	\$8,015,914	1	\$8,015,914		
Salem	2	\$118,616	1	\$8,223,090	0	\$0	0	0	2	\$118,616	1	\$8,223,090	3	\$8,341,705		
Somerset	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0		
Sussex	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0		
Union	0	\$0	1	\$1,238,150	0	\$0	0	0	0	\$0	1	\$1,238,150	1	\$1,238,150		
Warren	5	\$1,996,497	0	\$0	0	\$0	0	0	5	\$1,996,497	0	\$0	5	\$1,996,497		
Total	128	\$279,995,992	19	\$171,953,986	10	\$37,537,063	0	0	138	\$317,533,055	19	\$171,953,986	157	\$489,487,041		

Source: NJ OMB 2013; NJDEP 2013

Note: Total Value represents total structural and estimated content replacement value.

SFHA Special Flood Hazard Area



## Table 5.6-17. State Building Exposure to the 1% Annual Chance Flood Hazard by Agency

		A-7	Zone			V-Zone					_	SFHA		
		Owned		Leased	(	Owned	Lea			Owned	]	Leased		Total
Agency	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value
Agriculture	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0
Banking and Insurance	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0
Chief Executive	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0
Children and Families	1	\$3,655,514	5	\$50,078,015	1	\$887,897	0	0	2	\$4,543,410	5	\$50,078,015	7	\$54,621,425
Community Affairs	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0
Corrections	30	\$175,601,331	1	\$2,635,684	0	\$0	0	0	30	\$175,601,331	1	\$2,635,684	31	\$178,237,016
Education	0	\$0	1	\$35,841,320	0	\$0	0	0	0	\$0	1	\$35,841,320	1	\$35,841,320
Environmental Protection	55	\$14,772,836	2	\$6,876,559	8	\$36,254,497	0	0	63	\$51,027,334	2	\$6,876,559	65	\$57,903,893
Health	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0
Human Services	1	\$53,444,961	0	\$0	0	\$0	0	0	1	\$53,444,961	0	\$0	1	\$53,444,961
Judiciary	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0
Juvenile Justice Commission	7	\$1,840,888	0	\$0	0	\$0	0	0	7	\$1,840,888	0	\$0	7	\$1,840,888
Labor and Work Force Development	0	\$0	1	\$1,216,202	0	\$0	0	0	0	\$0	1	\$1,216,202	1	\$1,216,202
Law and Public Safety	0	\$0	1	\$25,833,976	0	\$0	0	0	0	\$0	1	\$25,833,976	1	\$25,833,976
Legislature	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0
Military and Veterans Affairs	12	\$16,335,540	1	\$451,732	0	\$0	0	0	12	\$16,335,540	1	\$451,732	13	\$16,787,273
Miscellaneous Commissions	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0



		A-7	Zone			V-Zone						SFHA		
	(	Owned		Leased	C	)wned	Lea	sed		Owned	J	Leased		Total
Agency	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value	Count	Total Value
Motor Vehicles Commission	1	\$4,962,410	3	\$18,781,004	0	\$0	0	0	1	\$4,962,410	3	\$18,781,004	4	\$23,743,415
Personnel	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0
State	0	\$0	0	\$0	0	\$0	0	0	0	\$0	0	\$0	0	\$0
State Police	6	\$4,137,151	3	\$7,416,815	1	\$394,669	0	0	7	\$4,531,821	3	\$7,416,815	10	\$11,948,636
Transportation	15	\$5,245,360	0	\$0	0	\$0	0	0	15	\$5,245,360	0	\$0	15	\$5,245,360
Treasury	0	\$0	1	\$22,822,677	0	\$0	0	0	0	\$0	1	\$22,822,677	1	\$22,822,677
Total	128	\$279,995,992	19	\$171,953,986	10	\$37,537,063	0	0	138	\$317,533,055	19	\$171,953,986	157	\$489,487,041

Source: NJ OMB 2013; NJDEP 2013c Note: Total Value represents total structural and estimated content replacement value.

SFHA Special Flood Hazard Area



County	Total Vulnerable	Airports	Special Needs	Communication	Correctional Institutions	Dams	Electric Power	EMS	EOCs	Ferries	Fire	Highway Bridges	Highway Tunnels	Light Rail Facilities	Medical	Military	Natural Gas	Oil	Police	Ports	Potable Water	Rail Facilities	Rail Tunnels	Schools	Shelters	Storage of Critical Records	Wastewater
Atlantic	109	0	1	0	0	36	1	12	0	0	14	9	0	0	1	0	0	0	6	0	1	0	0	19	8	0	1
Bergen	128	1	2	0	1	58	1	13	0	1	13	0	0	0	1	0	0	0	10	0	1	1	0	14	7	0	4
Burlington	168	0	0	0	0	130	0	1	0	0	4	3	0	2	0	0	0	0	2	0	0	0	0	7	10	0	9
Camden	62	0	0	0	1	44	0	1	0	0	1	3	0	2	0	0	0	0	1	1	1	0	0	2	4	0	1
Cape May	75	1	2	0	0	9	0	12	0	1	16	2	0	0	0	0	0	0	6	1	1	0	0	11	12	0	1
Cumberland	39	0	0	0	0	31	0	2	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	1
Essex	55	2	0	0	2	21	0	3	0	0	2	2	0	0	0	0	0	0	2	3	1	1	0	9	5	0	2
Gloucester	52	0	0	0	0	44	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	4	0	0	2
Hudson	112	0	2	1	1	0	2	6	0	8	11	3	1	12	5	0	1	0	2	3	1	6	1	30	12	0	4
Hunterdon	57	0	0	0	0	45	0	6	0	0	4	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0
Mercer	49	0	0	0	0	37	0	1	0	0	1	0	0	0	0	0	0	0	0	0	1	1	0	2	4	0	2
Middlesex	63	0	0	0	0	37	0	3	0	0	3	4	0	0	0	0	0	1	1	0	2	0	0	6	5	0	1
Monmouth	67	0	3	0	0	9	0	14	0	2	13	1	0	0	0	1	0	0	5	2	0	0	0	12	0	0	5
Morris	134	1	3	0	0	100	0	7	0	0	9	0	0	0	0	0	0	0	4	0	0	1	0	2	4	0	3
Ocean	135	0	1	0	0	68	0	22	0	0	17	4	0	0	0	0	0	0	12	0	0	1	0	9	1	0	0
Passaic	108	0	0	0	0	89	0	3	0	0	1	0	0	0	0	0	0	0	0	0	3	1	0	6	3	0	2
Salem	55	0	1	0	0	33	0	4	0	0	4	1	0	0	0	0	0	0	1	0	0	0	0	4	4	0	3
Somerset	86	0	1	0	0	55	0	7	0	0	3	0	0	0	0	0	0	0	2	0	1	3	0	5	8	0	1
Sussex	61	0	1	0	0	58	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
Union	61	0	2	0	0	25	1	4	0	0	2	1	0	0	0	0	0	0	3	5	1	0	0	11	4	0	2
Warren Total	44 1,707	0 5	0 19	0 1	0 5	28 <b>957</b>	0 4	5 125	0 0	0 12	1 122	4 38	0 1	0 16	0 7	0 1	0 1	0 1	0 56	0 11	0 14	0 15	0 1	2 154	3 <b>97</b>	0 0	1 44

# Table 5.6-18. State-Owned Critical Facilities in the 1% Annual Chance Flood Hazard Area (A- and V-Zones)

EMS Emergency Medical Service

EOC Emergency Operations Center



As the State of New Jersey continues to be developed, the state facilities will need to be located to conveniently serve the population base. As the New Jersey population continues to grow, so will the need for state services and facilities. Refer to the discussion earlier in this section regarding existing legislation and mitigation measures at the federal and state-level to reduce the impacts to future flood events.

## **Estimating Potential Losses by Jurisdiction**

Economic losses to New Jersey from flooding include but are not limited to: general building stock damage, agricultural losses, business interruption, impacts to tourism. These losses will negatively affect the tax base. Damage to general building stock can be quantified using HAZUS-MH as discussed above. Other economic components such as loss of facility use, functional downtime, and social economic factors are less quantifiable. For the purposes of this analysis, the general building stock damage is discussed further.

To estimate the potential losses by county, the HAZUS-MH flood model was used to estimate the potential losses to the general building stock. This analysis has been refined since the 2011 Plan due to the updated and improved flood hazard areas and depth grids across the State. Table 5.6-19 summarizes the estimated potential losses to the default general building stock by county. As statewide building data becomes available, the results will be updated to reflect more current data. The potential damage estimated to the general building stock inventory associated with the 1% annual chance flood is approximately \$33 billion which represents approximately 2.5% of the State's overall total general building stock inventory. Cape May county has the greatest estimated losses as a result of the flood event.

			SFHA
County	Total RCV	Estimated Loss	% of Total
Atlantic	\$38,043,171,000	\$2,829,243,000	7.4
Bergen	\$154,077,482,000	\$3,603,159,000	2.3
Burlington	\$62,700,794,000	\$897,247,000	1.4
Camden	\$70,467,051,000	\$739,741,000	1.0
Cape May	\$24,665,528,000	\$2,518,198,000	10.2
Cumberland *	\$18,128,613,000	\$199,891,000	1.1
Essex	\$113,124,687,000	\$1,778,606,000	1.6
Gloucester	\$33,534,660,000	\$352,724,000	1.1
Hudson	\$82,290,184,000	\$4,224,833,000	5.1
Hunterdon	\$21,720,513,000	\$302,380,000	1.4
Mercer	\$56,194,660,000	\$737,638,000	1.3
Middlesex	\$119,947,782,000	\$2,019,298,000	1.7
Monmouth	\$96,235,266,000	\$2,517,508,000	2.6
Morris	\$86,634,810,000	\$1,904,866,000	2.2

# Table 5.6-19. Estimated General Building Stock Losses from the 1% Annual Chance Flood Event, by County



			SFHA
County	Total RCV	Estimated Loss	% of Total
Ocean	\$73,559,915,000	\$4,665,078,000	6.4
Passaic	\$66,705,864,000	\$1,461,130,000	2.2
Salem *	\$8,092,037,000	\$342,894,000	4.2
Somerset	\$83,463,372,709	\$380,679,536	0.5
Sussex	\$20,979,595,000	\$110,198,000	0.5
Union	\$79,329,736,000	\$1,323,742,000	1.7
Warren	\$14,442,755,000	\$207,135,000	1.4
Total	\$1,324,338,475,709	\$33,116,188,536	2.5

Source: HAZUS-MH v 2.1

% percent

*RCV Replacement cost value. Total replacement cost value (structure and contents) as provided in the HAZUS-MH default general building stock inventory.* 

SFHA Special Flood Hazard Area

\*The estimated potential loss for Cumberland and Salem counties only represents loss within the areas associated with the preliminary work map depth grids provided by New Jersey Department of Environmental Protection (coastal areas); not loss in riverine flood hazard areas across the counties.

## **Estimating Potential Losses to State Facilities**

To estimate the potential loss to state facilities, the HAZUS-MH flood model and the statewide Land and Building Asset Management (LBAM) database provided by the NJ OMB were used. Direct building losses are the estimated costs to repair or replace the damage caused to the building. Tables 5.6-20 and 5.6-21 below summarize the estimated potential loss to state buildings by county and agency, respectively.

The potential damage estimated to state-owned and -leased buildings associated with the 1% annual chance flood is approximately \$73 million which represents approximately 1% of the total inventory. Hudson county has the greatest estimated potential loss as a result of the flood event. The New Jersey Department of Environmental Protection has the greatest estimated potential loss as a result of the flood event.



			Estimated Loss		
	Total RCV (structure and	Owned	Leased	Total	Percent of
County	contents)	RCV	RCV	RCV	Total
Atlantic	\$358,024,830	\$8,411,761	\$5,537,220	\$13,948,981	3.9
Bergen	\$219,423,769	\$197,830	\$0	\$197,830	0.1
Burlington	\$892,775,538	\$1,692,874	\$0	\$1,692,874	0.2
Camden	\$640,350,857	\$210,555	\$40,589	\$251,144	0.0
Cape May	\$117,950,706	\$447,975	\$43,454	\$491,429	0.4
Cumberland	\$813,708,672	\$622,786	\$0	\$622,786	0.1
Essex	\$674,467,788	\$20,964,377	\$1,464,319	\$22,428,696	3.3
Gloucester	\$76,531,777	\$0	\$0	\$0	0.0
Hudson	\$164,209,619	\$18,061,731	\$93,736	\$18,155,467	11.1
Hunterdon	\$411,264,979	\$127,786	\$0	\$127,786	0.0
Mercer	\$3,477,412,371	\$0	\$0	\$0	0.0
Middlesex	\$651,385,213	\$0	\$240,426	\$240,426	0.0
Monmouth	\$247,560,648	\$7,099,323	\$0	\$7,099,323	2.9
Morris	\$459,016,431	\$3,878	\$0	\$3,878	0
Ocean	\$172,110,712	\$1,377,176	\$206,023	\$1,583,199	< 1
Passaic	\$292,868,078	\$0	\$5,742,760	\$5,742,760	2
Salem	\$57,046,533	\$26,037	\$0	\$26,037	<1
Somerset	\$233,331,698	\$0	\$0	\$0	0
Sussex	\$49,168,422	\$0	\$0	\$0	0
Union	\$85,257,584	\$0	\$226,060	\$226,060	0.3
Warren	\$106,656,334	\$168,246	\$0	\$168,246	0.2
Total	\$10,200,522,559	\$59,412,335	\$13,594,587	\$73,006,922	<1

## Table 5.6-20. State Building Potential Loss to the 1% Annual Chance Flood Hazard, by County

Source: HAZUS-MH v2.1; NJOMB 2013

*RCV Replacement Cost Value. Total replacement cost value represents both structural value provided by the New Jersey Office of Management and Budget, and estimated contents.* 

			Estimated Loss									
	Total RCV (structure and	Owned	Leased	Total	Percent of							
Agency	contents)	RCV	RCV	RCV	Total							
Agriculture	\$2,876,615	0	\$0	\$0	0							
Banking and Insurance	\$83,777,640	0	\$0	\$0	0							
Chief Executive	\$12,653,376	0	\$0	\$0	0							
Children and Families	\$855,320,877	\$91,908	\$7,938,933	\$8,030,842	0.9							
Community Affairs	\$142,133,954	0	\$1,178,611	\$1,178,611	0.8							
Corrections	\$1,705,111,918	0	\$795,059	\$795,059	0.05							
Education	\$313,825,668	0	\$0	\$0	0							
Environmental Protection	\$466,946,331	\$25,004,726	\$0	\$25,004,726	5.4							
Health	\$146,433,703	0	\$0	\$0	0							
Human Services	\$1,689,928,602	0	\$0	\$0	0							
Judiciary	\$114,021,053	0	\$0	\$0	0							
Juvenile Justice Commission	\$258,880,851	\$517,754	\$0	\$517,754	0.2							
Labor and Work Force Development	\$242,663,875	0	\$2,442,543	\$2,442,543	1.0							
Law and Public Safety	\$498,665,653	0	\$646,230	\$646,230	0.1							
Legislature	\$165,085,389	0	\$0	\$0	0							
Military and Veterans Affairs	\$954,650,961	\$15,392,871	\$240,426	\$15,633,297	1.6							
Miscellaneous Commissions	\$15,650,656	0	\$0	\$0	0.0							
Motor Vehicles Commission	\$928,029,459	0	\$1,045,336	\$1,045,336	0.1							
Personnel	\$8,513,417	0	\$0	\$0	0							
State	\$208,816,705	0	\$0	\$0	0							
State Police	\$473,621,856	\$990,010	\$206,023	\$1,196,033	0.3							
Transportation	\$512,199,066	\$9,810,161	\$0	\$9,810,161	1.9							
Treasury	\$400,714,935	0	\$6,706,333	\$6,706,333	1.7							
Total	\$10,200,522,559	\$51,807,429	\$21,199,494	\$73,006,922	<1							

## Table 5.6-21. State Building Potential Loss to the 1% Annual Chance Flood Hazard by Agency

Source: HAZUS-MH v2.1; NJOMB 2013

*RCV* Replacement cost value. Total replacement cost value represents both structural value provided by the New Jersey Office of Management and Budget, and estimated contents.





The replacement cost values for critical facilities outside of the LBAM database were not available for this planning effort. As these data become available, the State will update this section of the Plan with new information.

Roads are the primary resource for evacuation to higher ground before and during the course of a riverine, coastal flood or tsunami event. Bridges exposed to flood events can be extremely vulnerable due to the forces transmitted by the wave run-up and by the impact of debris carried by the wave action. The forces of coastal flood and tsunami waves can also impact above ground utilities by knocking down power lines and radio/cellular communication towers. Power generation facilities can be severely impacted by both the velocity impact of the wave action and the inundation of floodwaters.

Flooding can cause extensive damage to public utilities and disrupt the delivery of services. Loss of power and communications may occur; and drinking water and wastewater treatment facilities may be temporarily out of operation. Flooded streets and roadblocks make it difficult for emergency vehicles to respond to calls for service. Floodwaters can wash out sections of roadway and bridges (Foster 2010).

The replacement cost values for critical facilities outside of LBAM were not available for this planning effort. For the purposes of this 2014 Plan update, the exposure analysis methodology was used to estimate potential losses to critical facilities and infrastructure. A total risk exposure would be equal to the full replacement value of each critical facility exposed.

## **Environmental Impacts**

Floods are naturally occurring events that benefit riparian systems that have not been disrupted by human development. These benefits include groundwater recharge and sediment movement that replenishes nutrients to agricultural soils. The shifting sediment keeps the elevation of a land mass above sea level. Floods can also lead to negative impacts on the environment. Loss of riparian buffers, land use change within a watershed, and introduction of non-natural contaminants may cause environmental issues when floods occur (Montz and Tobin 1997; Rubin 2013).

The basic environmental impact of major flooding is morphological; the shape of the river valley is often determined more by a catastrophic event. This process is a primary factor in forming the natural habitat for flora and fauna and may influence habitats beyond the river corridor. Therefore, floods are major, direct determinants of the natural environmental and, indirectly, of the human uses of the river corridor (Hickey and Salas 1995).

Flooding (inland, coastal, or tsunami waves) can cause a wide range of environmental impacts. These include, but are not limited to generating large amounts of tree and construction debris, dispersing household hazardous waste into the fluvial system, and contaminating water supplies and wildlife habitats with extremely toxic substances. Floods of greater depth are likely to result in greater environmental damage than floods of lesser depth. Long duration floods could exacerbate environmental problems because clean-up will likely be delayed and contaminants have the potential of remaining in the environment for a longer period of time. Cleaning up after a flood presents additional environmental concerns. The volume of debris to be collected, the extent to which public utilities (water supply systems and sewer operations) have been damaged, and the quantity of agricultural and industrial pollutants entering water bodies might present additional issues (Montz and Tobin 1997; Rubin 2013).